

Summary

The work presented in this thesis consists of synthesis, characterization and study of molecular properties of Schiff base coordinated polynuclear copper(II) complexes. Three different multidentate hydroxyl rich Schiff base ligands (E)-2-((1-hydroxybutan-2-ylimino)methyl)phenol, (E)-2-((1-hydroxybutan-2-ylimino)methyl)-6-methoxyphenol and 2-ethoxy-6-[(1-hydroxymethyl-propylimino)-methyl]-phenol used for the synthesis of complexes. The complexes are characterized by elemental analysis, IR, ESI-mass and X-ray single crystal structure determination. Variable temperature magnetic susceptibility measurements were carried out in the range 2-300 K. Interactions of complexes with the calf thymus DNA (CT-DNA) and serum albumins were investigated using electronic absorption and fluorescence spectroscopic techniques.

Chapter 1 contains general introduction and literature survey about 3d metal coordination compounds and their potential applications, such as magnetic, fluorescence and biological activity.

Chapter 2, describes synthesis and characterization of three novel tetranuclear copper(II) complexes namely $[\text{Cu}_4(\text{L}^1)_4] \cdot (\text{dmf})$ (**1**), $[\text{Cu}_4(\text{L}^1)_4]$ (**2**) and $[\text{Cu}_4(\text{L}^2)_2(\text{HL}^2)_2(\text{H}_2\text{O})_2] \cdot 2(\text{ClO}_4) \cdot 6(\text{H}_2\text{O})$ (**3**) (H_2L^1 , (E)-2-((1-hydroxybutan-2-ylimino)methyl)phenol; H_2L^2 , (E)-2-((1-hydroxybutan-2-ylimino)methyl)-6-methoxyphenol). The complexes were synthesised by the reaction of copper(II) perchlorate and the tridentate Schiff base ligands. Complexes **1** and **2** possess face-sharing dicubane core structure, on the other hand complex **3** has double open cubane core structure. Variable temperature magnetic susceptibility measurements in the range 2-300 K indicate an overall weak antiferromagnetic exchange coupling in all complexes.

Chapter 3, present two copper(II) complexes, $[\text{Cu}_4(\text{L})_2(\text{HL})_2(\text{H}_2\text{O})_2] \cdot 2(\text{ClO}_4) \cdot 2(\text{H}_2\text{O}) \cdot \text{DMF}$ (**1**) and $[\text{Cu}_4(\text{L})_2(\text{HL})_2(\text{H}_2\text{O})_2] \cdot (\text{tp})$ (**2**) [$\text{H}_2\text{L} = 2\text{-ethoxy-6-}[(1\text{-hydroxymethyl-propylimino)-methyl] \text{-phenol}$; $\text{tp} = \text{terephthalate}$] which were synthesized and characterized by single crystal X-ray diffraction and spectroscopic studies. The structural determination reveals that both the complexes are tetranuclear with double-open-cubane like core framework. Interactions of complexes with the calf thymus DNA (CT-DNA) and serum albumins were investigated using electronic absorption and fluorescence spectroscopic techniques.

Chapter 4, represent synthesis and characterization of Cu(II) complexes $[\text{Cu}_4(\text{L})_2(\text{HL})_2(\text{H}_2\text{O})_2] \cdot 2(\text{pv})$ (**1**) and $[\text{Cu}_4(\text{L})_2(\text{HL})_2(\text{H}_2\text{O})_2] \cdot (\text{ssal})$ (**2**) [$\text{H}_2\text{L} = 2\text{-Ethoxy-6-}[(1\text{-hydroxymethyl-propylimino)-methyl] \text{-phenol}$; $\text{pv} = \text{pivalate}$; $\text{ssal} = 2\text{-Hydroxy-5-sulfosalicylate}$]. Both the complexes are tetranuclear with double open cubane core framework. C-H... π interactions results the formation of 1D supramolecular structure for both the complexes. The interactions of complexes with the calf thymus DNA, bovine serum albumin (BSA) and human serum albumin (HSA) were studied using UV-vis absorption and fluorescence spectroscopic techniques. Molecular docking technique has been used to determine the mode of interaction of complexes with CT-DNA and serum albumins.

Chapter 5, reports the synthesis of two 1D polynuclear copper(II) compounds $\{[\text{Cu}_2(\text{HL}^1)_2(\text{fum})] \cdot (\text{H}_2\text{O}) \cdot (\text{MeOH})\}_n$ (**1**) and $\{[\text{Cu}_2(\text{L}^2)_2(\text{tp})] \cdot (\text{H}_2\text{O})\}_n$ (**2**) ($\text{fum} = \text{fumarate}$; $\text{tp} = \text{terephthalate}$) using alkoxo, hydroxo, phenoxo and imine donor Schiff bases $[(\text{E})\text{-2-}((1\text{-hydroxybutan-2-ylimino)methyl)phenol]$ (H_2L^1) and $[(\text{E})\text{-2-}((1\text{-hydroxybutan-2-ylimino)methyl)-6\text{-methoxyphenol}]$ (H_2L^2). Structure determination reveals that the complexes **1** and **2** crystallize in the monoclinic system with space group $P2_1/n$ and Pn , respectively. Variable temperature magnetic property study indicates overall antiferromagnetic exchange coupling in both the complexes. The interaction of complexes

with calf thymus DNA (CT-DNA) and bovine / human serum albumins were studied using electronic absorption and fluorescence spectroscopic techniques.

Chapter 6, contain synthesis and characterization of two copper(II) complexes namely $[\text{Cu}_4(\text{L}^1)_4] \cdot 0.5(\text{H}_2\text{O})$ (**1**) and $[\text{Cu}_4(\text{L}^2)_2(\text{HL}^2)_2(\text{H}_2\text{O})_2](\text{sq}) \cdot 2(\text{H}_2\text{O})$ (**2**) ($\text{H}_2\text{L}^1 = [(\text{E})\text{-}2\text{-}((1\text{-hydroxybutan-}2\text{-ylimino)methyl)phenol}]$, $\text{H}_2\text{L}^2 = [(\text{E})\text{-}2\text{-}((1\text{-hydroxybutan-}2\text{-ylimino)methyl)-6\text{-methoxyphenol}]$, sq = squarate and tp = terephthalate ion). The structural determination reveals that complex **1** crystallizes in the triclinic system with space group $\text{P}\bar{1}$, whereas complexes **2** crystallize in the monoclinic system with space group $\text{P}2_1/\text{n}$. Both **1** and **2** are tetranuclear species with Cu_4O_4 close cubane core and double-open cubane like core structure, respectively. Variable temperature magnetic property study indicates overall antiferromagnetic exchange coupling in both complexes. The PHI program was used to study their magnetic behavior. The Hamiltonians of type $H = -J_1 (S_1S_2 + S_1S_4 + S_2S_3 + S_3S_4) - J_2 (S_1S_3 + S_2S_4)$ and $H = -J_1 (S_1S_3 + S_2S_4) - J_2 (S_1S_2 + S_3S_4) - J_3S_1S_4$ were used to study magnetic behavior of **1** and **2**, respectively, and good agreement between the experimental and simulated results were found by using the parameters: $g = 2.14$, $J_1 = -20.2 \text{ cm}^{-1}$ and $J_2 = -1.7 \text{ cm}^{-1}$ for **1**, and $g_1 = 2.14$, $g_2 = 2.12$, $J_1 = -54.8 \text{ cm}^{-1}$, $J_2 = -15.9 \text{ cm}^{-1}$ and $J_3 = 0 \text{ cm}^{-1}$ for **2**.