




BIBLIOGRAPHY



BIBLIOGRAPHY

- Adams, M. L., Philpot, W. D. and Norvell, W. A. (1999). Yellowness index: an application of spectral second derivatives to estimate chlorosis of leaves in stressed vegetation. *Int. J. Remote Sens.*, 20(18), 3663–3675.
- Ahn, J. H., Hahm, Y. I., Park, C. S., Kim, B. H. and Kim, J. K. (1994). Modelling for prediction of potato late blight (*P. Infestans*) by meteorological elements. *RDA journal of agricultural Sciences Crop Protection*, 36(1), 320–325.
- Ahn, J. H., Hahm, Y. I., Park, E. W. (1998). Development of “moving average method” for prediction of initial appearance of potato late blight. *Korean Journal of Plant Pathology*, 14(1), 34–40.
- Andersson, B., Widmark, A. K. and Yuen, J. (2009). The role of oospores in the epidemiology of potato late blight. Proceedings of the III international Late Blight Conference, Beijing, China, April 3–6, 2008, *Acta Horticulture*, p.834.
- Andrade-Piedra, J. L., Forbes, G. A. and Shtienberg, D. (2005a). Qualification of a plant disease simulation model: performance of the LATEBLIGHT model across a broad range of environments. *Phytopathology*, 95, 1412–1422.
- Andrade-Piedra, J. L., Hijmans, R., Forbes, G. A., Fry, W. E., Nelson, R. J., (2005b). Simulation of potato late blight in the Andes. I: modification and parameterization of the LATEBLIGHT model. *Phytopathology*, 95, 1191–1199.
- Andrivon, D. (1995). Biology, ecology, and epidemiology of the potato late blight pathogen *Phytophthora infestans* in soil. *Phytopathology*, 85, 1053–1056.
- Anon. (1947). The measurement of potato blight. *Trans. Brit. Mycol. Soc.*, 31, 140–141.
- Anonymous. (2010). Final Report for Asia Pacific Network Project, 2010.
- Apan, A., Held, A., Phinn, S. and Markley, J. (2004). Detecting sugarcane ‘orange rust’ disease using EO-1 Hyperion hyperspectral imagery. *Int. J. Remote Sensing*, 25(2), 489–498.
- Arora, R. K. (1987). Some factors affecting potato tuber infection and loss in yield by late blight. *Indian Phytopathology*, 40, 226–231.
- Arora, R. K., Sharma, K. K. and Bombawale, O. M. (1999). Management of potato late blight by changing planting date in north-western plain. *Indian J. Mycol. Pl. Pathol.*, 29(3), 355–358.

- Aylor, D. E. (1990). The role of intermittent wind in the dispersal of fungal pathogens. *Annual Review of Phytopathology*, 28, 73–92.
- Badhwar, G. D. (1984). Automatic corn-soybean classification using Landsat MSS data: II. Early season crop proportion estimation. *Remote sensing of Environment*, 14, 31–37.
- Basu, A. (2009). Employing eco friendly potato disease management allows organic tropical Indian production systems to prosper. *Journal of Food and Agro-Industry, Special issue*, S80–S87.
- Basu, S. K. (1913). The late blight of potato. *Agric. J. of Bihar & Orissa*, 1, 142–149.
- Bauer, M. E. (1975). The role of remote sensing in determining the distribution and yield of crops. *Advances in Agronomy*, 27, 271–304.
- Beaumont, A. (1947). The dependence on the weather on the dates of outbreak of potato bligh epidemics. *Trans. Br. Mycol. Soc.*, 31, 45–53.
- Beckett, M., Daughtrey, M. and Fry, W. (2005). Temperature and leaf wetness requirements for pathogen establishment, incubation period, and sporulation of phytophthora infestans on petunia x hybrida. *Plant Disease.*, 89(9), 975–979.
- Bhattacharyya, S. K., Phadatare, S. G., Khanna, R. N., Srivastava, D. S., Singh, D. S. and Prasad, B. (1983). Efficiency of some fungicides in controlling late blight of potato in India. *Indian J. Agric. Sci.*, 53, 152–157.
- Bhattacharyya, S. K., Singh, B. P., Singh, P. H. and Ram, S. (1987). Retardation of potato late blight by fungicides with eradicant and protectant and protectant properties. *Indian J. Plant Pathol.*, 5, 169–177.
- Bianchi, R., Cavalli, R. M., Fiumi, L., Marino, C. M. and Pignatti, S. (1999). Airborne MIVIS hyperspectral imaging spectrometer over natural and anthropic areas. In: *Proceedings of the Fourth International Airborne Remote Sensing Conference and Exhibition/21st Canadian Symposium on Remote Sensing*, vol.1, Ottawa, Ont., Canada, pp. 337–344.
- Blackburn, G. A. and Steele, C. M. (1999). Towards the remote sensing of matorral vegetation physiology relationships between spectral reflectance, pigment, and biophysical characteristics of semiarid bush land canopies. *Remote Sens. Environ.*, 70(3), 278–292.
- Bock, C. H., Parker, P. E., Cook, A. Z. and Gottwald, T. R. (2008b). Characteristics of the perception of different severity measures of citrus canker and the relations between the various symptom types. *Plant Dis.*, 92, 927–939.

- Bombawale, O. M., Bhattacharyya, S. K. and Ravindran, C. D. (1991). Factors affecting potato late blight development in Punjab. *Indian Phytopath.*, 43, 319–324.
- Bonham, C. D. (1989). Measurements for Terrestrial Vegetation. Wiley, New York. pp. 352.
- Bourke, P. M. A. (1964). Emergence of potato late blight, 1843–1846. *Nature*, 203, 805–808.
- Bravo, C., Moshou, D., West, J., McCartney, A. and Ramon, H. (2003). Early disease detection in wheat fields using spectral reflectance. *Biosyst Eng.*, 84, 137–145.
- Brown, J. F., Reed, B. C., Hayes, M. J., Wilhite, D. A., Hubbard, K. (2002). A prototype drought monitoring system integrating climate and satellite data. Pecora 15/Land satellite Information IV/ISPRS commission I/FIEOS 2002 conference proceedings.
- Bruck, R. I., Fry, W. E., Apple, A. E. and Mundt, C. C. (1981). Effect of protectant fungicides on the development stages of *Phytophthora infestans* in potato foliage. *Phytopathology*, 71, 164–166.
- Butler, W. (1903). Potato diseases in India. *Agril ledger crop Dises. Pest Ser.*, 7, 87–124.
- Butler, E. J. (1918). Fungi and diseases in plant. *Thacker Spenk and Cooperation*, Calcutta.
- Carlisle, D. J., Cooke, L. R., Watson, S. and Brown, A. E. (2002). Folier aggressiveness of northern Irland isolates of *Phtophthora infestans* on detached leaflets of three potato cultivars. *Plant Pathology*, 51, 424–434.
- Carlson, Tobu N., and Riziley, David A. (1997). On the relation between NDVI, fractional vegetation cover, and leaf area index. *Remote Sensing of Environment*, 62, 241–252.
- Carter, G. A., Rebbek, J. and Percy, K. E. (1995). Leaf optical properties in *Liriodendron tulipifera* and *Pinus strobes* as influenced by increased atmospheric ozone and carbon dioxide. *Can. J. For. Res.*, 25, 407–412.
- Carter, G. A., Cibula, W. G. and Miller, R. L. (1996). Narrow-band reflectance imagery compared with thermal imagery for early detection of plant stress. *J. Plant Physiol.*, 148, 516–523.
- Carter, G. A. and Knapp, A. K. (2001). Leaf optical properties in higher plants: linking spectral characteristics to stress and chlorophyll concentration. *American Journal of Botany*, 88, 677–684.
- Castro, K. L., Sanchez-Azofeifa, G. A. (2008). Changes in spectral properties, chlorophyll content and internal mesophyll structure of senescing *Populus balsamifera* and *Populus tremuloides* leaves. *Sensors*, 8(1), 51–69.
- Castro, A. I. D., Jurado-Expósito, M., Gómez-Casero, M. T., López-Granados, F. (2012). Applying neural networks to hyperspectral and multispectral field data for discrimination of cruciferous weeds in winter crops. *Sci. World J.*, 8, 1–11.

- Ceccato, P., Flasse, S., Tarantola, S., Jacquemoud, S. and Gregoire, J. M. (2001). Detecting vegetation leaf water content using reflectance in the optical domain. *Remote Sens. Environ.*, 77, 22–33.
- Chappelle, E. W., Kim, M. S. and McMurtrey, J. E. (1992). Ratio analysis of reflectance spectra (RARS): An algorithm for the remote estimation of the concentrations of chlorophyll *a*, chlorophyll *b*, and carotenoids in soybean leaves. *Remote Sens. Environ.*, 39, 239–241.
- Chatterjee, P. B. (1997). Epidemics of diseases and insect pests in West Bengal with analysis into their causes and impacts (1945–1947). *Frontier of Plant Protection*, Department of Plant Protection, Biswa Bharati, 1, 1–12.
- Cheng, T., Rivard, B., Sánchez-Azofeifa, G. A., Feng, J. and Calvo-Polanco, M. (2010). Continuous wavelet analysis for the detection of green attack damage due to mountain pine beetle infestation. *Remote Sens. Environ.*, 114, 899–910.
- Chester, K. S. (1950). Plant disease losses: their appraisal and interpretation. *Plant Dis. Rep.*, Supplement 190–198(S193), 190–362.
- Clark, R.N. and Roush, L. (1984) Reflectance Spectroscopy Quantitative Analysis Techniques for Remote Sensing Applications. *Journal of Geophysical Research*, 89, 6329–6340.
- Cobb, N. A. (1892). Contribution to an economic knowledge of the Australian rusts (Uredinae). *Agric. Gazt.*, (NSW) 3, 60.
- CPRI. (1987). Annual Scientific Report. Central Potato Research Institute, Shimla, India. pp. 96–109.
- CPRI. (1999). Annual Scientific Report. Central Potato Research Institute, Shimla, India. pp. 70–77.
- Crosier, W. (1934). Studies in the biology of *Phytophthora infestans* (Mont.) de Bary. *Memoir–Cornell University Agricultural Experiment Station (USA)*, 155, 1–40.
- Cui, Y. (2010). Extraction of vegetation fraction based on the dimidiate pixel model and vegetation index transform plan. *Proceedings of SPIE*, 7752(1), 775202-1–775202-8.
- Cui, Y., Luo, Y., Wang, L. (2011). Extraction of vegetation fraction based on the dimidiate pixel model and vegetation index transform plan. In: *Proceedings PIAGENG 2010: Photonics and imaging for agricultural engineering*, SPIE, U.S.A., 7752, p 775202.
- Curran, P. J., Dungan, J. L. and Peterson, D. L. (2001). Estimating the foliar biochemical concentration of leaves with reflectance spectrometry: Testing the Kokaly and Clark methodologies. *Remote Sens. Environ.*, 76(3), 349–359.

- Dastur, J. F. (1915). Potato blight in India. *Mem. Dept. Agric. India. Bot. Ser.*, 7, 163–176.
- Dastur, J. F. (1917). Conditions influencing the distribution of potato blight in India. *Agric. J. India Spec. Ind. Congress*, 12, 90–96.
- Datt, B. (1999). A new reflectance index for remote sensing of chlorophyll content in higher plants: tests using eucalyptus leaves. *J. Plant Physiol.*, 154, 30–36.
- Dawson, T. P. and Curran, J. (1998). New technique for interpolating the reflectance red edge position. *Int. J. Remote Sens.*, 19(11), 2133–2139.
- De Bary, A. (1876). Research into the nature of the potato fungus, *Phytophthora infestans*. *J. Res. Agaric. Soc. Eng.*, 12, 239–269.
- De, B. K. and Basu, A. (2002). Three decades study on monitoring of late blight disease of potato in the plains of West Bengal. *Environment and Ecology*, 20(1), 216–218.
- Delalieux, S., Somers, B., Verstraeten, W. W., van Aardt, A. N. J., Keulemans, W. and Coppin, P. (2009). Hyperspectral indices to diagnose leaf biotic stress of apple plants, considering leaf phenology. *International Journal of Remote Sensing*, 30(8), 1887–1912.
- Dewelle, G. A. (1964). Forecasting crop infection by the potato blight fungus. *Meda. Vesh. K. Hed. Met.Inst.*, 82, 144.
- Dey, N. R. (1947). Cultivation and storage of potatoes in Bihar with special reference to the disease prevalent in the stores and the fields. *Allahabad Emg.*, 21, 177–204.
- Dutt, B. L. (1964). Late blight of potato in India, I. Distribution and blight period. *Indian Potato J.*, 6, 34–41.
- Dutt, B. L. (1979). Bacterial and fungal diseases of potatoes. ICAR, New Delhi, p.196.
- Dutt, B. L. (1980). Late blight of potato in KK Nambiar (ed) Proc. *Workshop on Phytophthora disease of tropical plants*. CPCRI, Kasargod, Kerala, 19–23 Sept. 1980, pp. 61–73.
- Dutta, S., Singh, S. K. and Panigrahy, S. (2014). Assessment of late blight induced diseased potato crops: A case study for West Bengal district using temporal AWiFS and MODIS Data. *Journal of the Indian Society of Remote Sensing*, 42(2), 353–361.
- Fahim, M. A., Hassanien, M. K. and Mostafa, M. H. (2003). Relationships between climatic conditions and potato late blight epidemic in Egypt during winter season 1999-2001. *Applied Ecology and Environmental Research*, 1(1/2), 159–172.
- Feng, W., Shen, W., He, L., Duan, J., Guo, B., Li, Y. (2016). Improved remote sensing detection of wheat powdery mildew using dual-green vegetation indices. *Precis. Agric.*, 17, 608–627.

- Feng, W., Wang, X. Y., Song, X., He, L., Wang, C. Y. and Guo, T. C. (2013). Hyperspectral estimation of canopy chlorophyll density in winter wheat under stress of powdery mildew. *J. Agr. Eng. Ers.*, 29, 114–123.
- Filella, I. and Penuelas, J. (1994). The Red Edge Position and Shape as Indicators of Plant Chlorophyll Content, Biomass and Hydric Status. *International Journal of Remote Sensing*, 15, 1459–1470.
- Fisher, J. I., Richardson, A. D., Mustard, J. F. (2007). Phenology model from surface meteorology does not capture satellite-based green up estimations. *Global Change Biology*, 13, 707–721.
- Flier, W. G. and Turkensteen, L. J. (1999). Floiar aggressiveness of Phtophthora infestans in three potato growing regions in the Nethrlands. *European Journal of Plant Pathology*, 105, 381–388.
- Forbes, D. A. and Jarvis, M. C. (1994). Host resistance for management of potato late blight. In: *Advances in Potato Pest Biology And Management* (Eds. Zehnder, G., Jansson, R., Raman, K.V.), St Paul Minnesota, American Phytopathological Society, 439–457.
- Franke, J. and Menz, G. (2007) Multi-temporal wheat disease detection by multi-spectral remote sensing. *Precision Agric.*, 8, 161–172.
- Fraser, R. N. (1998). Hyperspectral remote sensing of turbidity and chlorophyll a among Nebraska Sand Hills lakes. *Int. J. Remote Sens.*, 19(8), 1579–1589.
- Fry, W. E. (1975). Integrated effects of polygenic resistance and a protective fungicide on development of potato late blight. *Phytopathology*, 65, 908-911.
- Fry.W. E., Bruck, R. I. and Mundt, R. I. (1979). Retardation of potato late blight epidemics by fungicides with eradicant and protectant properties. *Plant Disease Reporter*, 63, 970–974.
- Galford, G. L., Mustard, J. F., Melillo, J., Gendrin, A., Cerri, C. C., Cerri, C. E. P. (2008). Wavelet analysis of MODIS time series to detect expansion and intensification of row-crop agriculture in Brazil. *Remote Sens. Environ.*, 112, 576–587.
- Gallo, K. P. and Flesch, T. K. (1989). Large-area crop monitoring with the NOAA AVHRR: Estimating the silking stage of corn development. *Remote Sensing of Environment*, 27, 73–80.
- Gao, B. (1996). NDWI—a normalized difference water index for remote sensing of vegetation liquid water from space. *Remote Sensing of Environment*, 58, 257 – 266.
- Gates, D. M., Keegan, H. J., Schleter, J. C. and Weidner, V. R. (1965). Spectral properties of plants. *Appl. Optics*, 4, 11–20.

- Gates, D. M. (1968). Transpiration and leaf temperature. *Ann. Rev. Plant Physiol.*, 19, 211–238.
- Gausman, H. W. (1974). Leaf reflectance of near-infrared. *Photogrammetric Engineering & Remote Sensing*, 40(2), 183–191.
- Gausman, H. W. (1977). Reflectance of leaf components, *Remote Sensing of Environment*, 6(1), 1–9.
- Gausman, H. W., Allen, W. A. and Escobar, D. E. (1974). Refractive-index of plant-cell walls. *Appl. Opt.*, 13, 109–111.
- Genc, H., Genc, L., Turhan, H., Smith, S. E., Nation, J. L. (2008). Vegetation indices as indicators of damage by the sunn pest (Hemiptera: Scutelleridae) to field grown wheat. *Afr. J. Biotechnol.*, 7, 173–180.
- Gisi, U. (1991). Synergism between fungicides for control of Phytophthora. See Ref. 57a, 361–372.
- Gitelson, A. A. and Merzlyak, M. N. (1994). Spectral reflectance changes associated with Autumn senescence of *Aesculus hippocastanum* L. and *Acer platanoides* L. leaves. Spectral Features and Relation to Chlorophyll Estimation. *Journal of Plant Physiology*, 143, 286–292.
- Gitelson, A. A. and Merzlyak, M. N. (1996). Signature analysis of leaf reflectance spectra: algorithm development for remote sensing of chlorophyll. *J. Plant Physiol.*, 148, 494–500.
- Gitelson, A. A., Kaufman, Y. J., Stark, R. and Rundquist, D. (2002). Novel algorithms for remote estimation of vegetation fraction. *Remote Sens. Environ.*, 80, 76– 87.
- Gitelson, A. A., Zur, Y., Chivkunova, O. B. and Merzlyak, M. N., (2002). Assessing carotenoid content in plant leaves with reflectance spectroscopy. *Photochem. Photobiol.* 75, 272–281.
- Godínez-Alvarez, H., Herrick, J. E., Mattocks, M., Toledo, D. and van Zee, J. (2009). Comparison of three vegetation monitoring methods: their relative utility for ecological assessment and monitoring. *Ecol. Indicat.*, 9(5), 1001– 1008.
- Gong, P., Pu, R. and Heald, R. C., (2002). Analysis of in situ hyperspectral data for nutrient estimation of giant sequoia. *International Journal of Remote Sensing*, 23(9), 1827–1850.
- Grenville-Briggs, L. J., Aurova, A. O., Bruce, C. R., Willams, A., Whisson, S. C., Brich, P. R. J. and Van West, P. (2005). Elevated amino acid biosynthesis in Phtophthora

- infestans during appressorium formation and potato infection. *Fungal General Biology*, 42, 244–256.
- Guan, J. and Nutter, F. W. Jr. (2003). Quantifying the intrarater repeatability and interrater reliability of visual disease and remote sensing assessment methods in the alfalfa foliar disease pathosystem. *Can. J. Plant Pathol.*, 25, 143–149.
- Guenther, K. P. (1990). Vegetation Stress Monitoring by Fluorescence. European Space Agency—Special Publication (ESA SP). *ESTEC*, Noordwijk, The Netherlands, pp. 135–142.
- Hansen, M. C. and Loveland, T. R. (2012). A review of large area monitoring of land cover change using Landsat data. *Remote Sens. Environ.*, 122, 66–74.
- Hardisky, M. A., Klemas, V. and Smart, R. M. (1983). The influence of soil salinity, growth form, and leaf moisture on the spectral radiance of *Spartina alterniflora* canopies. *Photogramm. Eng. Remote Sens.*, 49, 77 – 83.
- Harrison, J. G. (1992). Effects of the aerial environment on late blight of potato foliage—a review. *Plant Pathol.*, 41, 384–416.
- Henfling, J. W. (1987). Late blight of potato: *Phytophthora infestans*. *Technical Information Bulletin 4 (second edition revised)*, CIP, Lima Peru, p.22.
- Hill, M. J. and Donald, G. E. (2003). Estimation of Spatio-temporal Patterns of Agricultural Productivity in Fragmental Landscapes Using AVHRR NDVI Time Series. *Remote Sensing of Environment*, 84, 367–384.
- Hmimina, G., Dufrêne, E., Pontailleur, J. Y., Delpierre, N., Aubinet, M., Caquet, B. (2013). Evaluation of the potential of MODIS satellite data to predict vegetation phenology in different biomes: an investigation using ground-based NDVI measurements. *Remote Sensing of Environment*, 132, 145–158.
- Horler, D. N. H., Dockray, M. and Barber, J., (1983). The red edge of plant leaf reflectance. *International Journal of Remote Sensing*, 4(2), 273–288.
- Horsfall, J. G. and Cowling, E. B. (1978). Pathometry: the measurement of plant disease (pp 120–136) In: Plant Disease: An Advanced Treatise. Vol II. J. G. Horsfall and E. B. Cowling, (eds.). *Academic Press*, New York.
- Huang, B., Zhang, H., Song, H., Wang, J., Song, C., (2013). Unified fusion of remote-sensing imagery: generating simultaneously high-resolution synthetic spatial–temporal–spectral earth observations. *Remote Sens. Lett.*, 4, 561–569.

- Huang, C., Goward, S. N., Masek, J. G., Thomas, N., Zhu, Z., Vogelmann, J. E. (2010). An automated approach for reconstructing recent forest disturbance history using dense Landsat time series stacks. *Remote Sens. Environ.*, 114, 183–198.
- Huang, J., Liao, H., Zhu, Y., Sun, J., Sun, Q., Liu, X. (2012). Hyperspectral detection of rice damaged by rice leaf folder (*Cnaphalocrocis medinalis*). *Comput. Electron. Agric.*, 82, 100–107.
- Huang, W., Lamb, D. W., Niu, Z., Zhang, Y., Liu, L., Wang, J., (2007). Identification of yellow rust in wheat using in-situ spectral reflectance measurements and airborne hyperspectral imaging. *Precision Agriculture*, 8, 187–197.
- Huete, A. R., R. D. Jackson, and D. F. Post, (1985). Spectral response of a plant canopy with different soil backgrounds. *Remote Sensing of Environment*, 17(1), 37–53.
- Huete, A., Didan, K., Miura, T., Rodriguez, E. P., Gao, X., Ferreira, L. G. (2002). Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sens. Environ.*, 83, 195–213.
- Hunt, Jr., E. R. and Rock, R.N. (1989). Detection of changes in leaf water content using near- and middle-infrared reflectances. *Remote Sensing of Environment*, 30, 43–54.
- Islam, A. S., Bala, S. K. (2013). Assessment of Potato Phenological Characteristics Using MODIS-Derived NDVI and LAI Information. *GIScience & Remote Sensing*, 45(4), 454–470.
- James, C. (1947). Potato late blight scale. British Mycological Society Manual, pp.60.
- James, W. C. (1974). Assessment of plant disease losses. *Ann. Rev. Phytopath.*, 12, 27–48.
- Janetos, A. C. and Justice, C. O., (2000). Land cover global productivity: a measurement strategy for the NASA programme. *Int. J. Remote Sensing*, 21(6&7), 1491–1512.
- Jiang, Z., Huete, A. R., Chen, J., Chen, Y., Li, J., Yan, G. and Zhang, X. 2006. Analysis of NDVI and scaled difference vegetation index retrievals of vegetation fraction. *Remote Sensing of Environment*, 101(3), 366–378.
- Jing, X., Yao, W., Wang, J. and Song, X. (2010). A study on the relationship between dynamic change of vegetation Coverage and precipitation in Beijing's mountainous areas during the last 20 years. *Math. Comput. Modell.*, pp. 1–7.
- Jones, C. D., Jones, J. B., Lee, W. S. (2010). Diagnosis of bacterial spot of tomato using spectral signatures. *Computers and Electronics in Agriculture*, 74, 329–335.
- Judelson, H. S. and Blanco, F. A. (2005). The spores of *Phytophthora*; Weapons of the plant destroyer. *Natl. Rev.*, 3, 47–58.

- Kadam, V. C., Sarode, M. S., Bendre, N. J., Shingte, V. V., Khot, S. B. and Lokhande, S. B. (1974). Late blight of potato *Phytophthora infestans* (Mont.) de. Bary. *Current Science*, 43, 260.
- Kankwatsa, P., Hakiza, J. J., Olanya, M., Kidanemariam, H. M. and Adipala, E. (2003). Efficacy of different fungicide spray schedules for control of potato late blight in South-western Uganda. *Crop Prot.*, 22, 545–552.
- Karnieli, A. (2003). Natural vegetation phenology assessment by ground spectral measurements in two semi-arid environments. *Int. J. Biometeorol*, 47, 179–187.
- Khan, M. A.; Rashid, A., Ullah, O. and Iqbal, M. J. (2003). Control of Late Blight of Potato by Foliar Application of Fungicides. *International Journal of Agriculture & Biology*, 5(4), 540–542.
- Khanna, S., Palacios-Orueta, A., Whiting, M. L., Ustin, S. L., Riano, D., Litago, J. (2007). Development of angle index for soil moisture estimation, dry matter detection and land-cover discrimination. *Remote Sens Environ.*, 109, 154–165.
- Kirk, W. W., Abu-El Samen, F. M., Muhinyuza, J. B., Hammerschmidt, R., Douches, D. S., Thill, C. A., Groza, H. and Thompson, A. L. (2005). Evaluation of potato late blight management utilizing host plant resistance and reduced rates and frequencies of fungicide applications. *Crop Protection*, 24, 961–970.
- Knapp, A. K., Carter, G. A. (1998). Variability in leaf optical properties among 26 species from a broad range of habitats. *Am. J. Bot.*, 85(7), 940–946.
- Knipling, E. B. (1970). Physical and Physiological Basis for the Reflectance of Visible and Near-Infrared Radiation from Vegetation. *Remote Sensing of Environment*, 1, 155–159.
- Kobayashi, K. and Salam. M. U. (2000). Comparing simulated and measured values using mean squared deviation and its components. *Agron. J.*, 92: 345–352.
- Kobayashi, T., Kanda, E., Kitada, K., Ishiguro, K. And Torigoe, Y. (2001). Detection of rice panicle blast with multispectral radiometer and the potential of using airborne multispectral scanners. *Phytopathology*, 91, 316–323.
- Kokaly, R. F., Clark, R. N., (1999). Spectroscopic Determination of leaf biochemistry using Band-Depth analysis of absorption features and stepwise multiple linear regression. *Remote Sensing of Environment*, 67(3), 267–287.
- Kumar, L., Schmidt, K., Dury, S. and Skidmor, A. (2001). Imaging Spectrometry and Vegetation Science, in: *Imaging Spectrometry: Basic Principles and Prospective Applications*. *Kluwer Academic Publishers*, Dordrecht, 111–155.

- Lacey, J. (1965). The infectivity of soils containing *Phytophthora infestans*. *Annals of Applied Biology*, 59, 363–380.
- Lal, T. B. (1949). Occurrence of late blight in the plains of India. *Indian Phytopath.*, 1, 164–181.
- Larsole A., Muhammed, H. H., Stafford J. V. (2007). Measuring crop status using multivariate analysis of hyperspectral field reflectance with application to disease severity and plant density. *Precis. Agric.*, 8, 37–47.
- Lelong, C. C. D., Pinet, P. C., Poilvé, H. (1998). Hyperspectral imaging and stress mapping in agriculture: a case study on wheat in Beauce (France). *Remote Sens. Environ.*, 66(2), 179–191.
- Leprieur, C., Kerr, Y. H., Mastorchio, S. and Meunier, J. C. (2000). Monitoring vegetation cover across semi-arid regions: comparison of remote observations from various scales. *International Journal of Remote Sensing*, 21(2), 281–300.
- Li, F., Chen, W., Zeng, Y., Zhao, Q., Wu, B. (2014). Improving Estimates of Grassland Fractional Vegetation Cover Based on a Pixel Dichotomy Model: A Case Study in Inner Mongolia, China. *Remote Sens.*, 6, 4705–4722.
- Lichtenthaler, H. K., Lang, M., Sowinska, M., Heisel, F. and Miehe, J. A. (1996). Detection of vegetation stress via a new high resolution fluorescence imaging system. *J. Plant Physiol.*, 148, 599–612.
- Lindow, S. E. (1983). Estimating disease severity of single plants. *Phytopathology*, 73, 1576–1581.
- Lins, E. C., Belasque, J. Jr. And Marcassa, L. G. (2009). Detection of Citrus canker in citrus plants using laser induced fluorescence spectroscopy. *Precision Agric.*, 10, 319–330.
- Liu, H., Xu, L., Ding, J., Deng, X. (2010). A new angle-based spectral index and its application in drought monitoring. *2010 Symposium on Photonics & Optoelectronics*, Chengdu, p. 1–4.
- Liu, Z., Huang, J. and Tao, R. (2008). Characterizing and estimating fungal disease severity of rice brown spot with hyperspectral reflectance data. *Rice Sci.*, 15, 232–242.
- López, M. M., Bertolini, E., Olmos, A., Caruso, P., Gorris, M. T., Llop, P., Penyalver, R. and Cambra, M. (2003). Innovative tools for detection of plant pathogenic viruses and bacteria. *International Microbiology*, 6, 233–243.
- Lorenzen, B. and Jensen, A. (1989) Changes in leaf spectral properties induced in barley by cereal powdery mildew. *Remote Sensing of Environment*, 27(2), 201–209.

- Madden, L. V., Hughes, G. and van den Bosch, F. (2007). *The Study of Plant Disease Epidemics*. APS Press, St. Paul, MN.
- Mahlein, A. K., Oerke, E. C., Steiner, U., Dehne, H. W. (2012a). Recent advances in sensing plant diseases for precision crop protection. *Eur. J. Plant Pathol.*, 133, 197–209.
- Mahlein, A. K., Steiner, U., Hillnhütter, C., Dehne, H. W., Oerke, E. C. (2012b). Hyperspectral imaging for small-scale analysis of symptoms caused by different sugar beet diseases. *Plant Meth.*, 8, 3.
- Mahlein, A. K., Rumpf, T., Welke, P., Dehne, H. W., Plümer, L., Steiner, U., Oerke, E. C. (2013). Development of spectral indices for detecting and identifying plant diseases. *Remote Sens. Environ.*, 128, 21–30.
- Malcolmson, J. F. (1976). Assessment of field resistance to blight (*Phytophthora infestans*) in potatoes. *Transactions of the British Mycological Society*, 67, 321–325.
- Malenovsky, Z., Albrechtova, J., Lhotakova, Z., Zurita-Milla, R., Clevers, G. P. W. and Schaepman, M. E., (2006). Applicability of PROSPECT model for Norway spruce needles. *International Journal of Remote Sensing*, 27(24), 5315–5340.
- Malik, A. K., Post, J. J., Smith, L. P. and Austin Barke, P. M. (1955). The forecasting from weather date of late blight of potato and other plant disease and pests. *Irish Met. Service Dept. of Indust. And Commer.* Dublin, Ireland.
- Malthus, T. J. and Madeira, A. C. (1993). High resolution spectroradiometry: spectral reflectance of field bean leaves infected by *Botrytis fabae*. *Remote Sensing Environmental*, 45, 107–116.
- Martínez, B., Gilabert, M. A. (2009). Vegetation dynamics from NDVI time series analysis using the wavelet transform. *Remote Sens. Environ.*, 113, 1823–1842.
- Maxwell, S. K., Sylvester, K. M. (2012). Identification of ever-cropped land (1984–2010) using Landsat annual maximum NDVI image composites: Southwestern Kansas case study. *Remote Sens. Environ.*, 121, 186–195.
- McGuire, A. D., Melillo, J. M., Randenon, J. T., Parton, W. J., Heimam, M., Meier, R. A., Clein, J. S., Kicklighter, D. W. and Sauf, W. (2000). Modeling the effects of snowpack on heteronophic respiration across northern temperate and high latitude regions: comparison with measurements of atmospheric carbon dioxide in high latitudes. *Biogeochem*, 58(1), 91-114.
- Mirik, M., Michels, G. J. Jr., Kassymzhanova-Mirik, S., Elliott, N. C., Catana, V., Jones, D. B., Bowling, R. (2006). Using digital image analysis and spectral reflectance data to

- quantify damage by greenbug (Hemitera: Aphididae) in winter wheat. *Comput. Electron. Agric.*, 51, 86–98.
- Mirik, M., Jones, D. C., Price, J. A., Workneh, F., Ansley, R. J., Rush, C. M. (2011). Satellite remote sensing of wheat infected by wheat streak mosaic virus. *Plant. Dis.*, 95, 4–12.
- Mirik, M., Ansley, R. J., Price, J. A., Workneh, F., Rush, C. M. (2013). Remote monitoring of wheat streak mosaic progression using sub-pixel classification of Landsat 5 TM imagery for site specific disease management in winter wheat. *Adv. Remote Sens.*, 2, 16–28.
- Mizubuti, E. S. G. and Fry, W. E. (1998). Temperature effects on developmental stages of isolates from three clonal lineages of *Phytophthora infestans*. *Phytopathology*, **88**(8), 837–843.
- Moshou, D., Bravo, C., West, J., Wahlen, S., McCartney, A., Ramon, H. (2004). Automatic detection of 'yellow rust' in wheat using reflectance measurements and neural networks. *Comput. Electron. Agric.*, 44, 173–188.
- Moshou, D., Bravo, C., Oberti, R., West, J., Bodria, L., McCartney, A., Ramon, H. (2005). Plant disease detection based on data fusion of hyper-spectral and multi-spectral fluorescence imaging using Kohonen maps. *Real Time Imag. J.*, 11, 75–83.
- Moulin, S., Kergoat, L., Viovy, N., Dedieu, G. (1997). Global-scale assesement of vegetation phenology using NOAA/AVHRR satellite measurements. *Journal of Climate*, 10, 1154–1170.
- Muhammed, H. H. 2005. Hyperspectral crop reflectance data for characterising and estimating fungal disease severity in wheat. *Biosys. Eng.*, 91(1), 9–20.
- Muhammed, H., Larsolle, A. (2003). Feature-vector based analysis of hyperspectral crop reflectance data for discrimination quantification of fungal disease severity in wheat. *Biosystems Engineering*, 86(2), 125–134.
- Namanda, S., Olanya, O. M., Adipala, E., Hakiza, J. J., Bedewy, R. E., Baghsari, A. S. and Ewell, P. (2004). Fungicides application and host resistance for potato late blight management: benefits assessments from on-farm studies in S. W. Uganda. *Crop Protection*, 23(11), 1075–1083.
- Newell, L. C. and Tysdal, H. M. (1945). Numbering and note taking systems for use in improvement of forage crops. *J. Amer. Soc. Agron.*, 37, 736–749.
- Nilsson, H. E. (1985a). Remote Sensing of 2-row Barley Infected by Net Blotch Disease. Va ¨xtskyddsrap porter–Jordbruk, 34. Uppsala, Sweden: Swedish Agricultural University, SLU.

- Nita, M., Ellis, M. A. and Madden, L. V. (2003). Reliability and accuracy of visual estimation of Phomopsis leaf blight of strawberry. *Phytopathology*, 93, 995–1005.
- Nutter, F. W. Jr., Teng, P. S., and Shokes, F. M. (1991). Disease assessment terms and concepts. *Plant Dis.*, 75, 1187–1188.
- Nutter, F. W. Jr. and Schultz, P. M. (1995). Improving the accuracy and precision of disease assessments: selection of methods and use of computer-aided training programs. *Can. J. Plant Path.*, 17, 174–185.
- Okamoto, H., Murata, T., Kataoka, T., Hata, S. I. (2007). Plant classification for weed detection using hyperspectral imaging with wavelet analysis. *Weed Biol. Manage.*, 7, 31–37.
- Palacios-Orueta, A., Khanna, S., Litago, J., Whiting, S., Ustin, L. (2006). Assessment of NDVI and NDWI spectral indices using MODIS time series analysis and development of a new spectral index based on MODIS shortwave infrared bands. *Proceedings of the 1st International Conference of Remote Sensing and Geoinformation Processing Trier, Germany*, p. 207–209.
- Pan, Z., Huang, J., Zhou, Q., Wang, L., Cheng, Y., Zhang, H., Blackburn, G. A., Yan, J., Liu, J. (2015). Mapping crop phenology using NDVI time-series derived from HJ-1 A/B data. *International Journal of Applied Earth Observation and Geoinformation*, 34, 188–197.
- Parlevliet, J. E. (1979). Components of resistance that reduce the rate of epidemic development. *Annal Review of Phytopathology*, 17, 203–222.
- Peñuelas, J., Baret, S. and Filella, I. (1995). Semi-empirical indices to access carotenoids /chlorophyll ratio from leaves spectral reflectance. *Photosynthetica*, 31, 221–230.
- Phukan, S. N. and Baruah, C. K. (1994). Studies on the reaction behaviour of potato plants to infection by late blight fungus *Phytophthora infestans*. *Advances in Plant Sciences.*, 7(1), 18–23.
- Phukan, S. N. (1995). Effect of environmental factors on the development of late blight disease of potato in Assam. *Journal of Living World*, 2(1), 10–13.
- Piao, S., Fang, J. Y., Zhou, L. M., Ciais, P., Zhu, B. (2006). Variations in satellite-derived phenology in China's temperate vegetation. *Global Change Biol*, 12, 672–685.
- Ponzoni, F. J. and Goncalves, J. L. (1999). Spectral features associated with nitrogen, phosphorous and potassium deficiency in Eucalyptus saligna seedling leaves. *International Journal of Remote Sensing*, 20, 2249–2264.

- Purevdorj, J. T. S., Tateishi, R. and Ishiyama, T. (1998). Relationships between percent vegetation cover and vegetation indices. *Int. J. Remote Sens.*, 19(18), 3519–3535.
- Rao, A. N. S. and Veeresh, G. K. (1989). Estimation of yield loss due to late blight in rainfed potato. *Current Research– University of Agricultural Sciences (Bangalore)*, 18(11), 157–158.
- Ray, S. S., Das, G., Singh, J. P. and Panigrahy, S. (2006). Evaluation of hyperspectral indices for LAI estimation and discrimination of potato crop under different irrigation treatments. *International Journal of Remote Sensing*, 27, 5373–5387.
- Ray, S. S., Singh, J. P. and Panigrahy, S. (2010) Use of hyperspectral remote sensing data for crop stress detection: ground based studies. *Int. Arch. Photogramm. Rem. Sens. Spatial Inform. Sci.*, 38, 562–567.
- Ray, S. S., Jain, N., Arora, R. K., Chavan, S. and Panigrahi, S. (2011). Utility of Hyperspectral Data for Potato Late Blight Disease Detection. *Journal of Indian Society of Remote Sens.*, 39(2), 161–169.
- Reader, J. (2009). Potato: A History of the Propitious Esculent. *Yale University Press*, p.315.
- Reed, B. C., Brown, J. F., Vander Zee, D., Loveland, R. T., Merchant, J. W., Ohlen, D. O. (1994). Measuring Phenological Variability from Satellite Imagery. *Journal of Vegetation Science*, 5, 703–714.
- Rouse, J. W., Haas, R. H., Schell, J. A. and Deering, D. W. (1974). Monitoring vegetation systems in the Great Plains with ERTS. In: *S.C. Freden, editor, Third earth resources technology satellite-1 symposium*. Vol. 1: Technical presentations. NASA SP-351. Natl. Aeronautics and Space Admin., Washington, DC. p. 309–317.
- Rush, C. (2002). Prediction, detection and quantification of plant diseases with remote sensing technologies. Precision Agriculture Initiative for Texas High Plains. Annual Comprehensive Report.
- Sakamoto, T., Yokozawa, M., Toritani, H., Shibayama, M., Ishitsuka, N., Ohno, H. (2005). A crop phenology detection method using time-series MODIS data. *Remote Sens. Environ.*, 96, 366–374.
- Sakamoto, T., Van Nguyen, N., Ohno, H., Ishitsuka, N., Yokozawa, M. (2006). Spatio-temporal distribution of rice phenology and cropping systems in the Mekong Delta with special reference to the seasonal water flow of the Mekong and Bassac rivers. *Remote Sens. Environ*, 100, 1–16.
- Saltelli, A. (2002). Sensitivity Analysis for Importance Assessment. *Risk Analysis*, 22(3), 1–12.

- Saltelli, A., Ratto, M., Andres, T., Campolongo, F., Cariboni, J., Gatelli, D., Saisana, M. and Tarantola, S. (2008). *Global Sensitivity Analysis: The Primer*. John Wiley & Sons.
- Sanches, I. D., Filho, C. R. S, Kokaly, R. F. (2014). Spectroscopic remote sensing of plant stress at leaf and canopy levels using the chlorophyll 680 nm absorption feature with continuum removal. *ISPRS Journal of Photogrammetry and Remote Sensing*, 97, 111–122.
- Saucedo, C. O. M., Mayea, S. S., Fernandez, P. I. E. and Zumaquero, P. O. (2002). Influence of folier microclimate in the prognosis of late blight in potato crops. *Centrao Agricola*, 29(4), 92–94.
- Savitzky, A. and Golay, M. J. E. (1964). Smoothing and differentiation of data by simplified least squares procedures. *Anal. Chem.*, 36, 1627–1639.
- Schaber, J. and Badeck, F. W. (2003). Physiology-based phenology models for forest tree species in Germany. *International Journal of Biometeorology*, 47, 193–201.
- Schmidt, K. S., Skidmore, A. K. (2003). Spectral Discrimination of vegetation of vegetation types in a coastal wetland. *Remote Sens. Environ.*, 85(1), 92–108.
- Schtienberg, D., Raposo, R., Bergeron, S. N., Legard, D. E., Dyer, A. T. and Fry, W. E. (1994). Incorporation of cultivar resistancve in reduced-sprays strategy to suppress early and late blight on potato. *Plant Disease*, 78(1), 23–26.
- Schuler, R. T. (2002). Remote Sensing Experiences in Production Fields. (<http://alfi.soils.wisc.edu/extension/FAPM/2002proceedings/Schuler.pdf>).
- Schwartz, M. D., Reed, B. C., White, M. A. (2002). Assessing satellite start-of-season (SOS) measures in the conterminous USA. *International Journal of climatology*, 22(14), 1793–1805.
- Shekhawat, G. S. (1990). Potato Diseases and Pests and their Management in the Hills of Himanchal Pradesh. *J. Indian Potato Assoc.*, pp. 94–101.
- Shen, W. Y., Li, Y., Feng, W., Zhang, H. Y., Zhang, Y., Xie, Y. X. (2015b). Inversion model for severity of powdery mildew in wheat leaves based on factor analysis-BP neural network. *J. Agr. Eng. Ers.*, 31, 183–190.
- Sherwood, R. T., Berg, C. C., Hoover, M. R. and Zeiders, K. E. (1983). Illusions in visual assessment of *Stagonospora* Leaf Spot of orchardgrass. *Phytopathology*, 73, 173–177.
- Sheskin, D. J. (1997). *Handbook of Parametric and Non-Parametric Procedures*. CRC Press, Boca Raton, FL.
- Shokes, F. M., Berger, R. D., Smith, D. H. and Rasp, J. M. (1987). Reliability of disease assessment procedures: A case study with late spot of peanut. *Oleagineux*, 42, 245–251.

- Simons, M. D. (1972). Polygenic resistance to plant disease and its use in breeding resistant cultivars. *Journal of Environmental Quality*, 1, 232–240.
- Sindhuja, S., Ashish, M., Reha, E., Christina, D. (2010). A review of advanced techniques for detection of plant diseases. *Comp. Electr. Agric.*, 72, 1–13.
- Singh, B. P. and Shekhawat, G. S. (1999). Potato late blight in India. *Tech. Bull. No. 27 (Revised)*, Central Potato Research Institute, Simla, pp. 85.
- Singh, D. (1996). Fungicidal spray schedule for economical management of potato late blight in North Western Hills of India. *Indian J. Mycol. P. Pathol.*, 26(3), 252-255.
- Singh, J. P., Trehan, S. P., Upadhaya, N. C. and Lal. S. S. (2008). Potato Based Cropping systems. In: *Twenty Steps Towards Hidden Treasure* (eds.), Pandey, S.K. and Chakraborty, S.K. *Central Potato Research Institute, Shimla, India*, p.113.
- Slaton, M. R., Hunt, E. R., Jr., and Smith, W. K. (2001). Estimating near-infrared leaf reflectance from leaf structural characteristics. *American Journal of Botany*, 88, 278–284.
- Soudani, K., Maire, G., Dufrêne, E., François, C., Delpierre, N., Ulrich, E., Cecchini, S. (2008). Evaluation of the onset of green-up in temperate deciduous broadleaf forests derived from Moderate Resolution Imaging Spectroradiometer (MODIS) data. *Remote Sensing of Environment*, 112, 2643–2655.
- Spits, H., Evenhuis, B. and Schepers, H. (2007). Fungicide dose rate and cultivar resistance, results of five years of field experiments in the Netherlands. *Tenth Workshop of an European Network for development of an Integrated Control Strategy of potato late blight Bologna, Italy*.
- Srikantaiaya, M. (1962). Late blight of potatoes in Mysore State. *Indian Potato J.*, 4, 49–50.
- Srivastava, S. N. S. (1962). Epidemic of late blight of potato in South India. *Plant Protection Bull.*, 14, 26–28.
- Starodub, I. L., Gurevich, B. I. and Filippov, A. V. (1993). Forecasting of *Phytophthora Infestans* (Mont.) de Bary development on potato plants based on the analysis of meteorological conditions during vegetative growth. *Mikologiya-i-Fitopatologiya*, 27(1), 54–59.
- Steddom, K., Heidel, G., Jones, D. and Rush, C. M. (2003). Remote detection of rhizomania in sugar beets. *Phytopathology*, 93, 720–726.
- Steven, M. D., Malthus, T. J., Demetriades-Shah, T. H., Danson, F. M. and Clark, J. A. (1990). High resolution spectral indices for crop stress. In M. D. Steven & J. A. Clark (Eds.), *Applications of remote sensing in agriculture*, 209–227.

- Stilwell, A. R., Hein, G. L., Zygielbaum, A. I., Rundquist, D. C. (2013). Proximal sensing to detect symptoms associated with wheat curl mitevectored viruses. *Intern. J. Remote Sens.*, 34, 4951–4966.
- Strachan, I. B., Pattey, E. and Boisvert, J. B. (2002). Impact of nitrogen and environmental conditions on corn as detected by hyperspectral reflectance. *Remote Sens. Environ.*, 80, 213 – 224.
- Sugiura, R., Tsuda, S., Tamiya, S., Itoh, A., Nishiwaki, K., Murakami, N., Shibuya, Y., Hirafujia, M., Nuske, S. (2016). Field phenotyping system for the assessment of potato late blight resistance using RGB imagery from an unmanned aerial vehicle. *Biosystem Engineering*, 148, 1–10.
- Taranik, J. V., Mouat, D. A., Elvidge, C. D. (1993). Hyperspectral technology for geologic applications, better understanding of earth environment. *Int. Geosci. Remote Sens. Symp. (IGARSS)*, 2, 917–920.
- Taubenhaus, J. J., Ezekiel, W. N. and Neblette, C. B. (1929). Airplane photography in the study of cotton root rot. *Phytopathology*, 19, 1025–1029.
- Thakur, M., Chandla, V. K. and Thakur, Y. (2012). Evaluation of indigenous technology for the management of whitefly, *Trialeurodes vaporariorum* L. (Homoptera: Aleuroidadae) in potato. *Potato J.*, 39 (1), 57–61.
- Toler, R. W., Smith, B. D. and Harlan, J. C. (1981). Use of aerial color infrared photography to evaluate crop disease. *Plant Disease*, 65, 24–31.
- Tucker, C. J., Elgin, Jr. J. H., Mc, M. III. J.E. and Fan, C. J. (1979). Monitoring corn and soybean development with hand-held radiometer spectral data. *Remote Sens. Environ.*, 8, 237-248.
- Tucker, C. J., Slayback, D. A., Pinzon, J. E., Los, S. O., Myneni, R. B., Taylor, M. G. (2001). Higher northern latitude normalized difference vegetation index and growing season trends from 1982 to 1999. *International Journal of Biometeorology*, 45, 184–190.
- Ullstrup, A. J., Elliott, C. and Hopppe, P. E. (1945). Report of the committee on methods for reporting corn disease ratings. *Mimeographed*, Unnumbered Publication of the USDA, Division of Cereal Crops and Diseases. 5 pp.
- Umaerus, V. And Lihnell, D. (1976). A laboratory method for measuring the degree of attack by *Phytophthora infestans*. *Potato Research*, 19, 91–107.
- Umaerus, V., Umaerus M., Erjefalt, L. and Nilsson, B. A. (1983). Control of *Phytophthora* by host resistance: problems and progress. In: *Phytophthora: Its biology, taxonomy,*

- ecology and pathology*. [Eds. D.S. Erwin, S. Barnicki-Garcia and P.H. Tsao], APS, St Paul Minnesota. pp. 315–326.
- Van Everdingen, E. (1926). Het. verband tusschen de weergesteldhied en de aarolppelziekte, *Phytophthora infestans* (The relation between weather conditions and potato late blight, *Phytophthora infestans*). *Tijdchr. Plantenziekten*, 32, 129–140.
- Vanderplank, J. E. (1968). Disease resistance in plants. *Academic Press.*, NewYork. p.206.
- Vigier, B. J., Pattey, E. and Strachan, I. B. (2004). Narrowband Vegetation Indexes and Detection of Disease Damage in Soybeans. *IEEE. Geoscience and Remote Sensing Letters*. 1(4), 255–259.
- Viña, A., Gitelson, A. A., Rundquist, D. C., Keydan, G., Leavitt, B., Schepers, J. (2004). Remote sensing: monitoring maize (*Zea mays* L.) phenology with remote sensing. *Agronomy Journal*, 96, 1139–1147.
- Wang, H., Guo, J., Ma, Z. (2012). Monitoring wheat stripe rust using remote sensing technologies in China. In: Li D, Chen Y (eds) *Computer and computing technologies in agriculture*, V. Springer, 163–175.
- Wang, Q., and Tenhunen, J. D. (2004). Vegetation mapping with multitemporal NDVI in North Eastern China transect (NECT). *International Journal of Applied earth Observation and Geoinformation*, 6, 17–31.
- Wardlow, B. D., Kastens, J. H., Egbert, S. L. (2006). Using USDA crop progress data for the evaluation of greenup onset date calculated from MODIS 250-meter data. *Photogrammetric Engineering and Remote Sensing*, 72, 1225–1234.
- Wardlow, B. D., Egbert, S. L., Kastens, J. H. (2007). Analysis of time-series MODIS 250 m vegetation index data for crop classification in the US Central Great Plains. *Remote Sensing of Environment*, 108, 290–310.
- Wardlow, B. D., Egbert, S. L. (2008). Large-area crop mapping using time-series MODIS 250 m NDVI data: An assessment for the US Central Great Plains. *Remote Sensing of Environment*, 112, 1096–1116.
- Wellman, F. L. (1972). Tropical American Plant Diseases. *Scarecrow Press*, Metuchen, New Jersey.
- West, J. S., Bravo, C., Oberti, R., Lemaire, D., Moshou, D. And McCartney, H. A. (2003) The potential of optical canopy measurement for targeted control of field crop diseases. *Ann. Rev. Phytopathol.*, 41, 593–614.

- White, M. A., Nemani, R. R., Thornton, P. E., Running, S. W. (2002). Satellite evidence of phenological differences between urbanized and rural areas of the eastern United States Deciduous Broadleaf Forest. *Ecosystems*, 5, 260–277.
- White, M. A., de Beurs, K. M., Didan, K., Inouye, D. W., Richardson, A. D., Jensen, O. P., O’Keefe, J., Zhang, G., Nemani, R. R., van Leeuwen, W. J. D., Brown, J. F., de Wit, A., Schaepman, M., Lin, X., Dettinger, M., Bailey, A. S., Kimball, J., Schwartz, M. D., Baldocchi, D. D., Lee, J. T., Lauenroth, W. K. (2009). Inter comparison, interpretation, and assessment of spring phenology in North America estimated from remote sensing for 1982–2006. *Global Change Biol*, 15, 2335–2359.
- Willmott, C. J. (1982). Some Comments on the Evaluation of Model Performance. *Bulletin of the American Meteorological Society*, 63, 1309-1313.
- Wilson, T. and Felt, R. (1998). Hyperspectral remote sensing technology (HRST) program. *IEEE Aerosp. Appl. Conf. Proc.*, 5, 193–200.
- Wisler, G. C. and Duffus, J. E. (2000). A century of plant virus management in the Salinas Valley of California ‘East of Eden’. *Virus Research*, 71, 161–169.
- Woodcock, C. E. And Strahler, A. H. (1987). The factor of scale in remote sensing. *Remote Sens Environ.*, 21, 311–332.
- Wu, B., Miaomiao, L., Changzhen, Y., Weifeng, Z. (2004). Developing method of vegetation fraction estimation by remote sensing for soil loss equation. *Geoscience and Remote Sensing Symposium 2004. IGARSS '04. Proceedings, 2004 IEEE International*, 6, 4352–4355.
- Wu, W., Yang, P., Tang, H., Zhou, Q., Chen, Z., Shibasaki, R. (2010). Characterizing spatial patterns of phenology in cropland of China based on remotely sensed data. *Agric. Sci. China*, 9, 101–112.
- Xiao-Bing, L., Yun-Hao, C., Pei-Jun, S. and Jin, C. (2003). Detecting Vegetation Fractional Coverage of Typical Steppe in Northern China Based on Multi-scale Remotely Sensed Data. *Acta Botanica Sinica.*, 45(10): 1146–1156.
- Xin, J., Yu, Z., Van Leeuwen, L., Driessen, P. M. (2002). Mapping crop key phenological stages in the North China Plain using NOAA time series images. *International Journal of Applied Earth Observation and Geoinformation*, 4, 109–117.
- Yang, C. (2009). Assessment of the severity of bacterial leaf blight in rice using canopy hyperspectral reflectance. *Precision Agriculture*, 11, 61–81.
- Yang, C. M. and Cheng, C. H. (2001). Spectral characteristics of rice plants infested by brown planthoppers. *Proceedings of Natural Science*, 25(3), 180–186.

- Yang, C. M. and Chen, R. K. (2004). Modeling rice growth with hyperspectral reflectance data. *Crop Science*, 44, 1283–1290.
- Yang, C. M., Cheng, C. H., Chen, R. K., (2007). Changes in spectral characteristics of rice canopy infested with brown planthopper and leaffolder. *Crop Science*, 47, 329–335.
- Yoshida, K., Schuenemann, V. J., Cano, L. M., Pais, M., Mishra, B., Sharma, R., Lanz, C., Martin, F. N., Kamoun, S., Krause, J., Thines, M., Weigel, D. and Burbano, H. A. (2013). The rise and fall of the *Phytophthora infestans* lineage that triggered the Irish potato famine. *eLife Sciences Publications Ltd*, eLife2:e00731.
- Yuan, L., Huang, Y., Loraamm, R. W., Nie, C., Wang, J., Zhang, J. (2014). Spectral analysis of winter wheat leaves for detection and differentiation of diseases and insects. *Field Crop. Res.*, 156, 199–207.
- Zhang, H., Chen, J., Huang, B., Song, H., Li, Y. (2013). Reconstructing seasonal variation of Landsat vegetation index related to leaf area index by fusing with MODIS data. *IEEE J. Select. Topics Appl. Earth Observ. Remote Sens.*, 1, 1–11.
- Zhang, J. C., Pu, R. L., Huang, W. J., Yuan, L., Luo, J. H., Wang, J. H. (2012a). Using *in-situ* hyperspectral data for detecting and discriminating yellow rust disease from nutrient stresses. *Field Crop. Res.*, 134, 165–174.
- Zhang, J. C., Pu, R. L., Wang, J. H. and Huang, W. J. (2012b). Detecting powdery mildew of winter wheat using leaf level hyperspectral measurements. *Comput. Electron. Agr.*, 85, 13–23.
- Zhang, J. C., Pu, R. L., Yuan, L., Wang, J. H., Huang, W. J. and Yang, G. J. (2014). Monitoring powdery mildew of winter wheat by using moderate resolution multi-temporal satellite imagery. *PLoS ONE*, 9(4), e93107.
- Zhang, J., Pu, R., Loraamm, R. W., Yang, G., Wang, J. (2014). Comparison between wavelet spectral features and conventional spectral features in detecting yellow rust for winter wheat. *Comput. Electron. Agr.*, 100, 79–87.
- Zhang, M., Liu, X., O'Neill, M. (2002). Spectral Discrimination of *Phytophthora infestans* infection on tomatoes based on principal component and cluster analyses. *Int. J. Remote Sens.*, 23(6), 1095–1107.
- Zhang, M., Qin, Z., Liu, X. and Ustin, S. (2003). Detection of stress in tomatoes induced by late blight disease in California, USA, using hyperspectral remote sensing. *International Journal of Applied Earth Observation and Geoinformation*, 4(4), 295–310.

- Zhang, M. and Qin, Z. (2004). Spectral analysis of tomato late blight infections for remote sensing of tomato disease stress in California. *IEEE 2004 International Geosciences and Remote Sensing Symposium*. VI: 4091-4094.
- Zhang, X., Friedl, M. A., Schaaf, C. (2006). Global Vegetation Phenology from Moderate Resolution Imaging Spectroradiometer (MODIS): Evaluation of Global Patterns and Comparison with in situ Measurements. *Journal of Geo-physical Research*, 111, G04017.
- Zhang, X., Friedl, M. A., Schaaf, C. B., Strahler, A. H., Hodges, J. C. F., Gao, F., Reed, B. C., Huete, A. (2003). Monitoring vegetation phenology using MODIS. *Remote sensing of environment*, 84, 471–475.
- Zhang, X., Liao, C., Li, J. and Sun, Q. (2013). Fractional vegetation cover estimation in arid and semi-arid environments using HJ-1 satellite hyperspectral data. *International Journal of Applied Earth Observation and Geoinformation*. 21(1), 506–512.
- Zhang, X., Liao, C., Li, J. and Sun, Q. (2013). Fractional vegetation cover estimation in arid and semi-arid environments using HJ-1 satellite hyperspectral data. *Int. J. Appl. Earth Observ. Geoinf.*, 21, 206–512.
- Zhang, Z., Cao, K., Gui, X. and Yang, Z. (2004). Advance on epidemic and forecast of potato late blight in China. In: *Proceedings of the Fifth World Potato Congress*, Kunming, China, pp.187–189.