

Total Pages—8

PG/IVS/PHS/402.1&402.2

/25

**M. Sc. 4th Semester Examination, 2025**

**PHYSICS**

**PAPER — PHS-402.1 & 402.2**

*Full Marks : 50*

*Time : 2 hours*

**Answer all questions**

*The figures in the right hand margin indicate marks*

*Candidates are required to give their answers in their own words as far as practicable*

**PAPER — PHS-402.1**

**( Statistical Mechanics-II )**

**[Marks : 20]**

**1. Answer any two questions : 2 × 2**

- (a) Find an expression of total number of particles of thin film (2D) obeying B.E. statistics.

*( Turn Over )*

(b) For  $j$ th mode of vibration dispersion

$$\text{relation } \omega_j = \sqrt{\frac{4k}{m}} \sin\left(\frac{j\pi}{2N}\right)$$

Find the density of states.

(c) Write down an expression of photoelectric current density for  $\hbar\omega \gg \phi$ .

(d) Write down Fluctuation-dissipation theorem.

2. Answer any *two* questions : 4 × 2

(a) In Landau theory of diamagnetism, write down an expression of grand partition

function and show that  $\bar{N} = \frac{zV}{\lambda^3} \frac{x}{\sinh x}$

where  $x = \frac{\hbar e H}{2\pi K_B T m}$  and other symbols

have usual meanings.

- (b) In Bragg willian approximation show that Long range order parameter near  $T_c$ ,

$$L = \left[ \frac{3(T_c - T)}{T} \right]^{1/2}$$

- (c) Find a expression of photoelectric current density for  $\hbar\omega < \text{work function } (\phi)$ .

- (d) Prove that for degenerate electron gas pressure  $P = \frac{2}{5} n E_f(0)$ .

3. Answer any *one* question :

8 × 1

- (a) In ID using chain of  $N$  sites,

$$H = -J_1 \sum_{i=1}^{N-1} S_i S_{i+1} - J_2 \sum_{i=1}^{N-2} S_i S_{i+2}$$

Find the correlation function  $\langle S_i S_{i+1} \rangle$

for  $N \rightarrow \infty$ .

( 4 )

(b) For 3D gas of bosons, single particle energy

$$\epsilon(p, n) = \frac{|p|^2}{2m} + An,$$

Where A is a positive constant and  $n = -j$  to  $j$  is an integer. Find an expression of  $\bar{N}/V$  and the condition of B.E. condensation.

[ Internal Assessment — 05 Marks ]

**PAPER — PHS-402.2**

( Nuclear Physics )

[Marks : 20]

1. Answer any *two* questions : 2 × 2

(a) Show that the scattering angles of neutron in both L-system and C-system are almost same if the collision is against a very heavy nucleus.

( 5 )

- (b) Present graphically how Bohr-Wheeler used the liquid-drop model to explain the process of nuclear fission reaction.
- (c) How would the limitations of liquid-drop model be resolved in the single-particle nuclear shell model ?
- (d) Write the disadvantages of nuclear reactor as a neutron source.

2. Answer any *two* questions : 4 × 2

(a) (i) Find the spin, parity and magnetic-moment of  ${}_{16}\text{S}^{33}$  nuclide using nuclear shell model.

(ii) Show that the maximum loss of energy in a single collision of neutron with the moderator nucleus is proportional to the initial energy of the neutron. 2 + 2

( 6 )

- (b) (i) The compound nucleus can become de-excited either through  $\gamma$ -emission or by fission into nuclear fragments. The mean life-time of the compound nucleus is found to be  $4.7 \times 10^{-15}$  sec and the partial width for  $\gamma$ -emission is  $\Gamma_{\gamma} = 3.4 \times 10^{-2} eV$ . Find the partial fission width of  $\Gamma_{\gamma}$  for the emission of fragment  $\gamma$ .
- (ii) How can you establish that neutrons at low energy behave like a gas ? 2 + 2
- (c) Write a short note on a crystal monochromator for neutrons. 4
- (d) (i) In case of harmonic oscillator nuclear shell model, establish the relation between the harmonic oscillator frequency and the nuclear mass number.

( 7 )

(ii) What are the experimental evidences in supports of the nuclear shell structure ?

3 + 1

3. Answer any *one* question :

8 × 1

(a) (i) The neutron and proton in a deuteron nucleus interact through a square-well potential of width  $b = 1.9 \times 10^{-15} \text{ m}$  and depth  $V_0 = 40 \text{ MeV}$  in the  $l = 0$  state. Calculate the probability that the proton moves within the range of the neutron. Use the approximation that  $m_n = m_p = m$  and  $kb = \pi/2$ ,

where  $k = \sqrt{m(V_0 - \epsilon_d)}/\hbar$  and  $\epsilon_d$  is the binding energy of the deuteron. Also, find the mean-square radius of the deuteron nucleus.

(ii) How can you establish that neutrons at low energy behave like a gas ? 5 + 3

- (b) (i) Find the expression for threshold energy ( $E_{th}$ ) for an endoergic nuclear reaction. Hence, show that

$$(E_{th})_{\theta=0} = -Q \left( 1 + m_x / M_X \right)$$

Symbols have their usual meanings.

- (ii) In case of the scattering of thermal neutrons wave with matter, find the expression for the refractive index of matter.

5 + 3

[ Internal Assessment — 05 Marks ]