

14. REFERENCES

- Abouel-Yazeed, A.M.(2013). Maintaining Quality and Extending Shelf –Life of Tilapia Oreochromis Niloticus Fish during Storage at 4°C, pub.Journal of the arabian aquaculture society, Vol. 8 No 2 , pp. 293-305.
- Ackman, R.G., 1994. Seafood lipids, in sea food. Chemistry, Processing Technology and Quality. London: Chaoman & Hall.
- Alaiz, M., Navarro, J. L., Vioque ,E. and Giron,J.,1992. Amino acid analysis by high-performance liquid chromatography after derivatization with diethylethoxymethylenemalonate. *J. Chromat. Anal.* 591, pp. 181-186.
- Ali, S. and Chakraborty, T., 1992. Bangladesh Mitha Panir Amerudandi Prani (A book on Freshwater Invertebrates of Bangladesh). Bangla Academy, Dhaka, Bangladesh, 1st edition, pp. 208.
- Amerine, M.A., Pangborn, R.N. and Roessler, E.B.,1965. Principle of sensory evaluation of food,Academic Place New york. Pp. 349.
- Annandale, N., 1921. The aquatic and amphibious mollusca of Manipur. The prosobranchia. *Records of the Indian Museum* 22,pp. 538-564.
- Anonymous, 2001. Official Methods of Analyses. 16th Edn., The Association of Official Analytical Chemists, Washington, DC., USA.
- Ansell, A.D., 1972. Distribution, growth and seasonal changes in biochemical composition for the bivalve *Donax littatus* _Da Costa. from Kames Bay, Millport, J. *Exp. Mar. Biol. Ecol.*, 10, pp. 137–150.

- Ansell, A.D.,1972. Seasonal changes in biochemical composition of the bivalve Chlamys septemradiata from the Clyde sea area,Marine Biol., 25,pp. 85-99.
- AOCS, 1991,Official Methods and Recommended Practice of American Oil Chemists' Society. 4th ed. Firestone,D. (Ed.). Washington, DC: Author.
- APHA,1992. Compendium of Methods for the Microbiological Examination of Food, 3rd Edn., (Vanderzant, C. and Splittstoesser,D, eds.) PHA, Washington DC.
- Ashton, I.P., 2002. Understanding lipid oxidation in fish. In: H.A. Bremner, editor. Safety and quality issues in fish processing. Cambridge, UK: Woodhead, Publishing, 3, pp. 254-286.
- Babu, A., Venkatesan, V.,Rajagopal , S., 2011.Fatty acid and amino acid compositions of *Thegastropods, tonnadolium* (linnaeus, 1758) and *Phalium glaucum* (linnaeus, 1758) from the gulf of mannar, southeast coast of india,pub Annals. Food Science and Technology, pp.159-163.
- Baby, R. L., Hasan, I., Kabir, K. A. and Naser, M. N., 2010, Nutrient Analysis of Some Commercially Important Molluscs of Bangladesh. pub.:Journal of Scientific Research;j.Sci.Res.2 (2), pp. 390-396.
- Banye B.L. 1976. Some effect of stress in the adult on the larval development of *Mytilus edulis*. Nature, 273, pp. 459.
- Bath, K.S., Kaur, H. and Dhillon, S.S.,1999.Correlation of molluscs with physico chemical factors at Harike Reservoir (Punjab). Indian J environ Sci. 3: 159-163.
- Beatty, S.A. and Gibbons, N.E., 1937. The measurement of spoilage in fish. Pub.:Journal of Biology Biodiversity Canada, pp. 77-91.
- Bender, D., 1992. Nutritional Biochemistry of the Vitamins. New York: Cambridge University Press, pp. 269-317.

- Beninger P.G. and Lucas A., 1984. Seasonal variations in condition, reproductive activity and gross biochemical composition of two species of adult clam reared in a common habitat: *Tapes decussatus L.* (Jeffreys) and *Tapes philippinarum* (Adams and Reeve), *J Exp Mar Biol Ecol*, 79, pp. 19-37.
- Brandt, R.A.M. 1974. The non-marine aquatic Mollusca of Thailand. In: Archiv für Molluskenkunde der Senckenbergischen Naturforschenden Gesellschaft, vol. 105. Senckenbergische Naturforschende Gesellschaft, pp.33.
- Brenna, J.T. 2002. Efficiency of conversion alpha-linolenic acid to long chain n-3 fatty acids in man. *Curr. Opin. Clin. Nutr. Metab. Care*, 5,pp. 127-132.
- Browne, A.A., 1978. Growth, mortality, fecundity, biomass and productivity of four lake populations of the prosobranch snail, *Viviparus georgianus*. *Ecology* 59(4). 742 - 750.
- Budha, P.B., Dutta, J. and Daniel, B.A., 2010. *Bellamya bengalensis*. The IUCN Red List of Threatened Species. Version 2014.2. (Web source : www.iucnredlist.org).
- Burdge, G.C., Jones, A.E. & Wootton, S.A. 2002. Eicosapentaenoic and docosapentaenoic acids are the principal products of ??-linolenic acid metabolism in young women. *Br. J. Nutr.*, 88,pp. 355-363.
- Cabuk, Y., Arsalan, N. and Yilmaz, V., 2004. Species composition and seasonal variations of the gastropoda in upper sakarya river system (Turkey) in relation to water quality. *Acta Hydrochimica et Hydrobiologic* 32 (6), pp. 393-400.
- Cagiltay,F., Erkan, N. Tosun, D. and Selcuk, A., 2011. Amino acid fatty acid, vitamin, and mineral contents of the edible garden snail (*Helix aspersa*). *Journal of Fisheries Sciences*, 5 (4), pp. 354-363.

- Calder, P.C. 2006. Polyunsaturated fatty acids and inflammation. Pros. Leuk. EFA, 75, pp. 197-202.
- Carrillo-Trip, M. & Feller, S.E. 2005. Evidence for a mechanism by which ?-3 polyunsaturated lipids may affect membrane protein function. Biochem., 44, pp. 10164-10169.
- Celek, P., 2014. Comparative study on biochemical composition of some edible molluscs at Canakkale coast, Turkey;International journal of Geo-Merine Sciences vol.43 (4), pp. 601-606.
- Chacko.D, Renitta.R.E., Patterson. J, 2005. Development of soup powder from squid Sepioteuthis lessoniana and shelf-lifeassessment during storage in laminated packaging material, pub Journal of food technology, pp. 449-452.
- Chakraborty, D., Mukherjee, M., Maity, J., 2014a. Seasonal Variation and Profitable Market Potentiality of *Bellamya bengalensis* in the Western Zone of West Bengal, India. Asian Resonance; vol.-III, issue-III, July, pp. 86-90.
- Chakraborty, D. and Mukherjee, M., 2014b. Livelihood Prospect of *Bellamya bengalensis*, its Proximate Protein and Amino Acid Composition & Culture in Captive Condition. Souvenir of National Brain Storming Workshop on Prospect , Potentials & Diversification of Culture System in Beel Fisheries Sector of North East India, Aquaculture & Biodiversity Centre, Dept. of Zoology, Guwahati University (Assam), Oct., pp. 12-24.
- Chakraborty, D., Mukherjee, M. and Maity, J., 2015a. Ethno-medicinal importance of viviparous gastropod, *Bellamya bengalensis* (lamarck, 1822) in terms of estimation of

- its proximate protein and amino acid composition. International Journal of Scientific Research, Volume : 4 | Issue : 1 | ISSN No 2277 – 8179, pp. 178-180.
- Chakraborty, D., Mukherjee, M. and Maity, J.,2015b. Estimation of proximate fatty acid composition in ethno-medicinally important viviparous gastropod, *Bellamya bengalensis* (lamarck, 1822) , International journal of advanced scientific and technical research ISSN 2249-9954, pub.Issue 5 volume 1, January-February 2015, pp. 84-92.
- Chakraborty, D., Mukherjee, M. and Maity, J.,2015c. Evaluation of market potentiality of *Bellamya bengalensis* at selected market areas in Purulia district, West Bengal, India.J.Environ. and Sociobiology: 12 (1), pp. 1-6
- Chen C. and Sung K., 1975. A preliminary study on reproduction and growth of the snail,*Bellamya aeruginosa* (veeve), Acta Hydrobiologica Sinica, cnki:ISSN:1000 3207.0-04-008.
- Choubisa, S. L. and Sheikh, Z.,2013. Freshwater snails (mollusca: gastropoda) as bioindicatorfor diverse ecological aquatic habitats, Cibtech Journal of Zoology ISSN: 2319–3883 (Online), Vol. 2(3) September-December, pp.22-26.
- Chourpagar, A.R. and Kulkarni, G.K., 2011. Effect of heavy metal pesticides on biochemical constituents in freshwater crab *Barytelphusa cunicularis* (westwood). Proc. Zool. Soc. India. 10(2), pp. 21-30.
- Chua, M.E., Sio, M.C., Sorongon, M.C., and Morales, M.L.,2013 The relevance of serum levels of long chain omega-3 polyunsaturated fatty acids and prostate cancer risk: a meta-analysis. Canadian Urological Association Journal 7 (5–6): E333–E343. doi:10.5489/cuaj.1056. PMC 3668400.PMID 23766835.

- Clarke, A.H., 1979. The fresh water molluscs of the Canadian interior basin, *Malacologia* 13, pp.1-509.
- Clemente, G., Mancini, M., Nazzaro, F., Lasorella, G., Rivieccio, A., Palumbo, A.M., Rivelles, A.A., Ferrara, L. and Giacco, R., 2003. Effects of different dairy products on postprandial lipemia. *Nutr. Metab. Cardiovasc. Dis.*, 13,pp. 377-383.
- Connell, J.J., 1975. Control of Fish Quality, (1st Edition). Fishing News Books Limited, London.
- Dabrowski, K., 1977. Protein requirements of grass carp fry (*Ctenopharyngodon idella* Val.). *Aquaculture* J., 12, pp.63–73.
- Dare P.J. and Edwards D.B., 1975. Seasonal changes in flesh weight and biochemical composition of mussels, *Mytilus edulis* _L., in the Conwy Estuary, North Wales, J. Exp. Mar. Biol. Ecol., 18, pp. 89–97.
- Delgado-Lista, J., Perez-Martinez, P., Lopez-Miranda, J., Perez-Jimenez, F. 2012. "Long chain omega-3 fatty acids and cardiovascular disease: a systematic review". The British journal of nutrition. 107 Suppl 2: S201–13.doi:10.1017/S0007114512001596. PMID 22591894,.
- De Zwaan, A. and Zandee, D.I., 1972. Body distribution and seasonal changes in the glycogen content of the common sea mussel, *Mytilus edulis* L. Comp. Biochem, Physiol, 43A, pp. 53–58.
- Diwan, A. D. and Nagabhushnam, R., 1970. Reproductive cycle and biochemical changes in the gonads of the freshwater crab, *Barytelphusa cunicularis* (westwood, 1836), pp. 164-173

- D.Menno, S., Peter G. and Anton J.W.,2009. *Viviparus acerosus* (Bourguignat, 1862) (Gastropoda: Viviparidae), a new exotic snail species for the Dutch fauna: Aquatic Invasions (2009) Volume 4, Issue 2, pp. 373-375.
- Dolson.L.,2014. List of High-Protein Foods and Amount of Protein in Each, Protein - High-Protein Foods and Amount of Protein in Each Food, About.com.
- Fahy, E., Subramanium, S., Brown, A.H., Glass, C.K., Merril Jr., A.H., Murphy, R.C., Raetz, C.R.H., Russell, D.W., Seyama, Y., Shaw, W., Shimizu, T., Spener, F., van Meer, G., VanNieuwenhze, M.S., White, S.H., Witztum, J.L. & Dennis, E.A. 2005. A comprehensive classification system for lipids. *J. Lipid Res.*, 46,pp 839-861.
- Falkner,G.,(1989, *Viviparus ater* am deutschen Bodensee- Ufer. *Heldia* 1(5/6), pp.188-189.
- Falniowski, A., Fialkowski, W., Szarowska, M. and Mazan, K., 1998. Shell biometry characters in species discrimination and classification within the genus *Viviparus* (Gastropoda: Architaenioglossa: Viviparidae) *Malakologische Abhandlungen*, vol 19, pp 29-45.
- FAO/WHO/UNU, 2007. "Protein and amino acid requirements in huma nutrition",WHO Press.http://whqlibdoc.who.int/trs/WHO_TRS_935_eng.pdf, page 150.
- Farahnak, A., Mobedi, I., Ghobadi, H. 2006; Nematode association with *Bellamya bengalensis* snail and its medical and veterinary importance in Khouzestanprovince (south west of Iran), Archives of Razi Institute, Vol. 61, No. 1, Spring (2006) 49-52.

- FAO/WHO, 2001. Human Vitamin and Mineral Requirements, pub.: Food and Nutrition Division, FAO Rome: pp 1-118.
- Folch, J., Lees, M. and Sloane-Stanley, G.H., 1957. A simple method for isolation and purification of total lipids from animal tissues. *J. Biol. Chem.*, 226, pp. 497-509.
- Francois, C.A., Connor, S.L., Bolewicz, L.C. & Connor, W.E. 2003. Supplementing lactating women with flaxseed oil does not increase docosahexaenoic acid in their milk. *Am. J. Clin. Nutr.*, 77, pp. 226-233.
- Gabbott P.A., 1975. Storage cycles in marine bivalve molluscs: A hypothesis concerning the relationship between glycogen metabolism and gametogenesis. In: Barnes, H. Ed., Proc. 9th Europ. Mar. Biol. Symp., Aberdeen Univ. Press, pp. 191-211.
- Gabbot P.A. and Bayne, B.L., 1973. Biochemical effects of temperature and nutritive sites of *Mytilus edulis* (L.). *Journal Marine Biological Association of the United Kingdom*, 53, pp. 269-286.
- Galtsoff, P.S., 1964. The American oyster (*Crassostrea virginica*). *United States Fish and Wildlife Service's Fisheries Bulletin*, 64, 480.
- Garbott, P.A., 1983. Developmental and seasonal metabolic activities in marine molluscs. In the *Mollusca. Envi. Biochem. and Physiology*, P.W. Hochachka, Academic Press, New York, 2, pp. 165-217.
- Ghabshi Aliya Al, Humaid Al-Khadhuri, Nasser Al-Aboudi, Sami Al-Gharabi, Aziz Al-Khatiri, Nashwa Al-Mazrooei and Ponnerassery Sukumaran Sudheesh, 2012. Effect of the Freshness of Starting Material on the Final Product Quality of Dried Salted Shark, *Advance Journal of Food Science and Technology* 4(2), pp. 60-63.

- Ghafoorunissa, A., Vanni, R., Laxmi, R. & Sesikeran, B. 2002. Effects of dietary alpha-linolenic acid from blended oils on biochemical indices of coronary heart diseases in Indians. *Lipids*, 37,pp. 1007-1086.
- Ghobadi, H. and Farahnak, A., 2004. A Faunistic Survey on the Cercariae of *Bellamya (Viviparus) bengalensis* Snails and Their Zoonotic Importance: Iranian J Publ Health, Vol. 33, No. 2, pp.38-42.
- Glöer, P., 2002. Die Süßwassergastropoden Nord- und Mitteleuropas. Bestimmungsschlüssel, Lebensweise, Verbreitung. Die Tierwelt Deutschlands 73 Teil. Conchbooks, Hackenheim.
- Goel, S.C., 1999. Principles of animal developmental biology, Himalayan Publishing House, Girgaon, Bombay, India, pp. 400-004.
- Gongongopadhyay, P.L., 2001. Fundamental of Business Management. Pub: Dey Book Concern, pp.1-673.
- Gopakumar, K., 2002. Biochemical Composition of Fish. Textbook of Fish Processing Technology, (Directorate of Information and Publishing of Agriculture ICAR, New Delhi), pp.18-30.
- Gopal,T.K.S., Thankamma, R. Shenoy, A.V. Rao, C.V.N. and Govindan, T.K., 1988. Development of flexible packaging for fish soup powder. In : The First Indian Fisheries Forum, Proceeding, (Joseph, M.M. Ed.), Asian Fisheries Society, Indian Branch, Mangalore, pp. 369-372.
- Government of India Planning Commission (GOIPC), 2013. Press Note on Poverty Estimates, 2011-12. Government of India press information bureau, New Delhi: 22 July, 2013, pp. 1-9.

- Goyens, P.L.L.,2006. Conversion of ?-linolenic acid in humans is influenced by the absolute amounts of ?-linolenic acid and linoleic acid in the diet and not by the ratio. Am. J. Clin. Nutr., 84, pp.44-35.
- Goyens, P.L.L., Spiker, M.E., Zock, P.L., Katan, M.B. & Mensink, R.P. 2006. Conversion of alpha-linolenic acid in humans is influenced by the absolute amounts of alpha-linolenic acid and linoleic acid in the diet and not by their ratio,(<http://www.ajcn.org/cgi/content/abstract/84/1/44>).American Journal of Clinical Nutrition 84 (1), pp. 44–53.
- Gupta, S. D., 1976. Macrofauna of Loni reservoir,J. Ind. Fish. Soc.India 8: 49-59.
- Gupta, P.C., Khatta, V.K. and Mandal, A.B., 1992. Analytical techniques in animal nutrition. Pub-Directorate of publications, Chaudhury Charan Singh, Haryana Agricultural University, Hisar-125 004,pp. 1-100.
- Haldar, A., Dey, T. K., Dhar, P.1 and Chakrabarti, J.,2014. Exploring the Nutritive Values of the Fresh Water Mussel *Lamellidens marginalis* as Potential Functional Food, IOSR Journal of Environmental Science, Toxicology and Food Technology (IOSR-JESTFT) e-ISSN: 2319-2402,p- ISSN: 2319-2399. Volume 8, Issue 8 Ver. III (Aug. 2014), pp. 01-07.
- Hamdani, A, and Mazouni, S. N., 2011. Changes in Biochemical Composition of the Gonads of *Donax trunculus* L. (Mollusca, Bivalvia) from the Gulf of Annaba (Algeria) in Relation to Reproductive Events and Pollution; Jordan Journal of Biological Sciences; Volume 4, Number 3, September, pp 149 – 156.
- Hedge, J.E. and Hofreiter, B.T., 1962. In: Carbohydrate Chemistry, 17 (Eds. Whistler R.L. and Be Miller, J.N.), Academic Press, New York.

- Heinze, V.M., and Actis A.B., 2012. Dietary conjugated linoleic acid and long-chain n-3 fatty acids in mammary and prostate cancer protection: a review. International journal of food sciences and nutrition 63 (1), pp. 66–78.
- Hongwei, Yan. Qi Li., Wenguang, Liu., Ruihai, Yu. and Lingfeng, K., 2009. Seasonal changes in reproductive activity and biochemical composition of the razor clam *Sinonovacula constricta*. Marine Biology Research. 6(1), pp.78-88.
- Imelda F. Pagulayan and Joel C. Cepillo (1990), The Reproductive Biology and Laboratory Maintenance of *Vivipara angularis* Muller (Prosobranchia:Viviparidae) : Transactions of the National Academy of Science and Technology,pp. 535-568.
- IS Food Safety, 2012, Manual method of analysis of foods, Meat and meet product Fish and fishery product, pub> Govt. of India, New Delhi, 2012,pp 31-33.
- Iyer, T.S.G., Kandoran, M.K., Thomas, M. and Mathew, P.T., 2000. Text book of quality assurance in sea food procession. Pub. Central Institute of Fisheries Technology and Society of Fisheries Technologists, pp 1-230
- John, G., 1980. Studies on *Anadra rhombea* (Born) from Porto-Novo coastal waters, Ph.D., thesis, Annamalai University, 190.
- Jump, D.B. 2008. N-3 polyunsaturated fatty acid regulation of hepatic gene transcription. Curr. Opin. in Lipidol., 19: 242-247.
- Kaneniwa, M., S. Miao, C., H. Iida and Y. Fukuda, 2000. Lipid components and enzymatic hydrolysis of lipids in muscle of Chinese freshwater fish. Journal of the American Oil Chemists' Society, 77: 825-830.
- Karina, A. Rodrigues, G. J. M., Massa, A., María and Militelli, I., 2013. Seasonal analysis of condition, biochemical and bioenergetic indices of females of Brazilian

- flathead, *Percophis brasiliensis*; *Neotropical Ichthyology*; Neotrop. ichthyol. vol.11 no.1 Porto Alegre Jan./Mar.
- Khan, R.A. and Chaudhuri, S., 1984. The population and production ecology of a freshwater snail, *Bellamya bengalensis* (Lamarck) (Gastropoda : Viviparidae) in an artificial lake of calcutta, India. Bull. Zool. Surv. India, 5 (2& 3): 59-76.
- Kotwal, S., Jun, M., Sullivan, D., Perkovic, V. and Neal, B. (2012). "Omega 3 Fatty Acids and Cardiovascular Outcomes: Systematic Review and Meta-Analysis". *Circ Cardiovasc Qual Outcomes*, 5 (6): 808–18.doi:10.1161/CIRCOUTCOMES.112.966168.PMID 23110790.
- Krishnamoorthi, K.P. and Sarkar, R., 1979. Macro- invertebrates as indicators of water quality. Proc. Symp. Environ. Biology. Academy of Environmental Biology, Muzaffarnagar, India. S.R. Verma, A. K. Tyagi and S.K. Bansal (eds.) pp. 133-138.
- Kumar, A. and Gupta, R.S., 1984. Anode on the sensitivity of enterotoxigenic *Staphylococcus aureus* for garlic extract (*Allium sativum*). Ind Vet. J., 61: 718-719.
- Labuze, T.P., 1970. Oxidative changes in foods at low and intermediate moisture levels, In, Duckworth, R.B. (ed.), water relations of Foods. Academic Press, New York, pp. 455-474.
- Lakshmanan, P.T..2000. Quality assurance in sea food processing; pub. CIFT and SFT (India), pp. 36-38.
- Lee, S.Y., 1986. Growth and reproduction of the greenmussel *Perna viridis* (L.) (Bivalvia: Mytiliacea) in contrasting environments in Hong Kong. Asian Marine Biol., 3, PP. 111-127.
- Liza, D. and Castle F., 2010. Omega-3 and Omega-6 Fatty Acids, pub by Neb Guide; <http://extension.unl.edu/publications,>

- Lodeiros, C.J., Rengel, J.J., Guderley, H.E., Nuseni, O. and Himmelman, J.H., 2001. Biochemical composition and energy allocation in the tropical scallop *Lyropecten (Nodipecten) nodosus* during the months leading up to and following the development of gonads, *Aquaculture*. 199, pp. 63-72.
- Love, R.M., 1992. Biochemical dynamics and the quality of fresh and frozen fish. *Fish Processing Technology*, Editors: Hall GM, (Chapman and Hall), pp 1-30.
- Ma, T., Gong, S., Zhou, K., Zhu, C., Deng, K., Luo, Q. and Wang, Z., 2010. Laboratory culture of the freshwater benthic gastropod *Bellamya aeruginosa* (Reeve) and its utility as a test species for sediment toxicity; *J Environ Sci (China)*. 2010;22(2), pp 304-13.
- MacDonald B.A. and Thompson R.J., 1986. Influence of temperature and food availability on the ecological energetics of the giant scallop *Placopecten magellanicus*, *Mar. Biol.*, 93,pp. 37-48.
- Mahata, M.C., 2002. Edible Shell Fish (Molluscs) of Chotanagpur Plateau, Jharkhand (India). Baripada, Orissa: Bio-publications, pp. 1-133 .
- Mahata, M.C., 2012. Freshwater edible molluscan species diversity and their sustainable development in the state of Jharkhand : review. *Biodiversity for Human Welfare*, pp 186-193.
- Malathi, S. and Thippeswamy, S., 2013. the proximate and mineral compositions of freshwater mussel *parreysia corrugata* (mullar, 1774) from tunga river in the western ghats, India. *G.J.B.A.H.S.*, Vol.2(3):165-170.
- Masniyom, P., Soottawat, B., & Visessanguan, W., 2005. Combination effect of phosphate and modified atmosphere on quality and shelf-life extension of

- refrigerated seabass slices. Journal of Food Science and Technology, 38, pp.745–756.
- Masalha, R., Chudakov, B., Muhamad, M., Rudoy, I., Volkov, I. and Wirguin I, 2001. Cobalamin-responsive psychosis as the sole manifestation of vitamin B₁₂ deficiency. Israeli Medical Association Journal 3, pp. 701–703. PMID 11574992.
- Masniyom, P., Soottawat, B. and Visessanguan, W. 2005. Combination effect of phosphate andmodified atmosphere on quality and shelf-life extension of refrigerated seabass slices. Journal of Food Science and Technology, vol.38, pp. 745–756.
- Michael, R. G., 1968. Studies on the bottom fauna in a tropical freshwater fish pond. Hydrobiologia 31,pp. 203-230.
- Misra K. K., Shkrob, I., Rakshit, S. and Dembitsky, V. M., 2001. Variability in fatty acids and fatty aldehydes in different organs of two prosobranch gastropod mollusks, Published by Elsevier Science,Biochemcal systematic & ecology,vol 30, Pages 749-761.
- Mitra and Dey,1992. Land and Fresh Water Molluscs (Part-I):Pub Handbook of Freshwater Molluscs of India. Calcutta: Zoological Survey of India.pp1-51.
- Monzon, A.B., 1985. Snails of medical importance in Family Viviparidae of Thailand. Paper read at the Training Course in Applied Malacology held in Bangkok, Thailand, November-December,1985.
- Morais, S., Boaventura, D. Narciso, Lui 's , Re Pedro, Hawkins S. J., 2003.Gonad development and fatty acid composition of *Patella depressa* Pennant (Gastropoda:Prosobranchia) populations with different patterns of spatial

- distribution, in exposed and sheltered sites; Journal of Experimental Marine Biology and Ecology,294; pp. 61–80.
- Mozaffarian, D., Ascherio, A., Hu, F., Stampfer, M.J., Willett, W.C., Siscovick, D.S. Rimm, E.B., 2005. Interplay between different polyunsaturated fatty acids and risk of coronary heart disease in men. Circulation, 111, pp. 157-164.
- Morton B., 1978. The population dynamics of *Anomalocardia squamosa* Lamarck (Bivalvia: Veneracea) in Hong Kong, J. Moll. Stud. 44, pp. 135-144.
- Mothet, J.P., Parent, A.T., Wolosker, H., Brady, R.O., Linden, D.J., Ferris, C.D., Pogawski, M. A., and Snyder, S.H. 2000. PNAS, 97, pp. 4926-4931.
- Mukherjee, M. and Basu, S., 2008. Commercially cultured of fresh water snail & edible oyster (in Bengali version). Pub: Dept. of Fisheries, Govt. of West Bengal. pp 1-67.
- Mukhopadhyay, T. and Ghosh .S, 2003. Lipid profile and fatty acid composition in egg of common carp (*Cyprinus carpio*),. J.Oleo Scence 52:439-442.
- Nagabhushanam R. and Mane, U.H., 1978. Seasonal variation inthe biochemical composition of *Mytilus viridis* at Ratnagiri, on west coast of India. Hydrobiologia, 57,pp. 69-72 .
- Nesemann H, S Sharma, G Sharma, SN Khanal, B Pradhan, DN Shah and RD Tachamo. 2007. Aquatic invertebrates of the Ganga. 14 Himalayan journal of sciences vol 7 issue 9 2011 Research paper river system:, Volume 1: Mollusca, Annelida, Crustacea (in part), pp. 263.
- Nordsieck, R., 2012. Amazing facts on molluscs, The homepage on gastropods, bivalves and other molluscs by Robert Nordsieck. Support and scripting: Martina Eleveld, 2351 Wiener Neudorf Austria (Web source : <http://www.molluscs.at/index.html>).

- Nordsieck, R., 2012. Fresh Water Snails, Part 1: Snails Breathing With Gills, 2351 WienerNeudorfAustria.(Websource:<http://www.molluscs.at/gastropoda/index.html?/gastropoda/freshwater.html>)
- Odum, E.P.1989. Input management of production system. Science 234, pp.117-181.
- Office of Dietary Supplements, National Institutes of Health (ODSNIH), 2011. Dietary Supplement Fact Sheet: Vitamin B12.. Retrieved 28 September.
- Pagulayan, I. F. 1986. Comparative characterization of the foot muscle, hemolymph, hepatopancreas proteins of *Radix quadrasi* and *Mixas cumingiana* (Pulmonata: Lymnaeidae).Unpublish ed material.
- Pagulayan, F.I. and Cepillo,C.J.,1990. The Reproductive Biology and Laboratory Maintenance of Vivipara angularis Muller (Prosobranchia:Viviparidae), Pub. Transactions of the National Academy of Science and Technology, pp. 535-567.
- Pal, M., Mukhopadhyay, T., and Ghosh, S., 2011. Proximate Fatty Acid and Amino Acid Composition of Fish Muscle and Egg Tissue of *Hilsa (Tenualosa ilisha)*, pub: Journal of Aquatic Food Product Technology, Taylor & Francis 20, pp. 160-171.
- Pardeshi, P. B. and Vedpathak A. N.,2013. Biochemical composition of freshwater bivalve, *Lamellidens marginalis*, (Lamark) from lentic and lotic water environments of Nathsagar reservoir at Paithan; Deccan current science; DCSI 09: pp. 198 – 204.
- Patterson, J.,2004. Smoking an ideal method to preserve Mollusk meat; SPS Trochus information, Bulletin #11 – September.
- Paul, D. J. (2003): Sustainable America's Aquatic Biodiversity Fresh Snail Biodiversity & Conservation; Publication Number 420-530, Posted October.

- Pawar, P.P., 2011. Preparation of battered and breaded product from freshwater fish (*Catla catla*),M.F.Sc thesis submitted to Konkan Krishi Vidyapeeth, Dapoli, Maharashtra state, India.
- Periyasamy N, Srinivasan M, Devanathan K, Balakrishnan S. (2011), Nutritional value of gastropod *Babylonia spirata* (Linnaeus, 1758) from Thazhanguda, Southeast coast of India;pub: Asian Pacific Journal of Tropical Biomedicine,pp S249-S252.
- Peterson, C.H. and Fegley, S.R., 1986. Seasonal allocation of resources to growth of shell, soma, and gonads in *Mercenaria mercenaria*, Biol. Bull., 171, pp. 597-610.
- Pharmacy and Therapeutics,2008. Omega-3-acid Ethyl Esters (Lovaza) For Severe Hypertriglyceridemia 22 Canadian Foo,.
- Pir, J., Tali, I., Mudgal, L. K. and Seddique, A., 2010. Distribution of Molluscans in Narmada River, India: sciencepub : 2(10), pp. 41-46.
- Pravakar, A.K. and Roy,S.P.,2009. Ethno-medicinal uses of some shell fishes by people of Kosi river basin of north-Bihar. India: Ethno-Med, 3(1),pp. 1-4.
- Preston, H.B., 1915. The fauna of British India including Ceylon and Burma : Mollusca (Freshwater Gastropoda and Pelecypoda). Taylor and Francis, Lond., pp.1-244.
- Qingzhu,2003. Quality indicators of northern shrimp (*pandalus borealis*) stored under different cooling conditions. Pub UNU-Fisheries Training Programme, pp. 1-83.
- Ramakrishna and Dey, A. 2007. Handbook on Indian freshwater molluscs. Zoological Survey of India, Kolkata.
- Ramon, M., Abello, P. and Richardson, C.A., 1995. Population structure and growth of *Donor trunculus* (Bivalvia: Donacidae) in the western Mediterranean. Mar. Biol., 121, pp. 665-671.

- Rathod,N. and Pagarkar, A., 2013. Biochemical and sensory quality changes of fish cutlets, made from pangasius fish (*Pangasianodon hypophthalmus*), during storage in refrigerated display unit at -15 to -18°C, pub. International Journal of Food, Agriculture and Veterinary Sciences, pp. 1-8.
- Ratnayake, W.M.N., Hansen S.L. & Kennedy, M.P. (2006). Evaluation of the CP-Sil 88 and SP-2560 GC columns used in the recently approved AOCS Official Method Ce 1h-05: Determination of cis-, trans-, saturated, monounsaturated and polyunsaturated.
- Rao, K. 1969. Mollusca have many uses. *Indian Farming*, 29(9): 41-45 and 62.
- Reddy, M.A., Elavarasan, A., Reddy, D.A. and Bhandary, M.H.,2012. Suitability of reef cod (*Epinephelus diacanthus*) minced meat for preparation of ready to serve product. Advances in Applied Science Research 3(3), pp 1513-1517.
- Reeds, P.J., 2000. Dispensable and indispensable amino acids for humans". J. Nutr. 130 (7): 1835S–40S.
- Roy, S.P.,2003. Secondary productivity of fresh water ecosystem. In: JS Datta Munshi, J Ojha, TKGhosh (Eds.): *Advances in Fish Research*. Delhi: Narendra Publishing House, Volume 3: pp. 99-108.
- Roy, S.P., Singh, B.K. , 2007. Ethno-medicinal uses of animal diversity by tribal people of Santhal Pargana in Jharkhand, India, In: AP Das, AK Pandey (Eds.): *Advances in Ethno-botany*. Dehra Dun: Bishen Singh Mahendra Pal Singh, pp. 211-217.
- Saha B.K., Jahan M.S. and Hossain M.A., 2013, Taxonomic record and distribution pattern of the banded pond snail, *Bellamya bengalensis* (Lamarck) (Gastropoda : Mesogastropoda) from pond water habitat of Rajshahi. Bangladesh J. Sci. Ind. Res. 48(1) pp 71-74.

- Sallam, K.I., 2007. Chemical, sensory and shelf-life evaluation of sliced salmon treated with salts of organic acids, Published in final edited form as: Food Chem. 2007 ; 101(2), pp 592–600.
- Saritha, K. and Patterson, J., 2014. Innovative Seafood Health Mix Powder from Juveniles of *Leiognathus* sp. and Their Quality for Human Consumption; pub. World Journal of Dairy & Food Sciences 9 (2), pp 113-120.
- Saudari Pawiro,2005, Bivalves:global production & trade; INFOFISH Internationl 1/, pp.45.
- Schileyko, A.A. & A.G. Kuznetsov. 1996. A new genus of the Subulinidae (Pulmonata) from Nepal. *Ruthenica* 5(2), pp.158-160.
- Sethi, N.K., Robilotti, E. and Sadan, Y.,2005. Neurological Manifestations Of Vitamin B₁₂ Deficiency. The Internet Journal of Nutrition and Wellness 2 (1). doi:10.5580/5a9.
- Shigmates, H. and Takeshita, H.,1959. On the changes in the weight of fat body and its chief constituents in the silkworm, *Bombyx mori L.* during metamorphosis. Appl. Entm. Zool. Japan. 3, pp. 123-126
- Shafakatullah, N. Shetty, S., Lobo, R.O. and Krishnamoorthy, M., 2013. Nutritional Analysis of Freshwater bivalves, *Lamellidens* spp. from River Tunga, Karnataka, India; Research Journal of Recent Sciences, Vol. 2 (ISC-2012), pp.120-123 .
- Shafakatullah, N. and Krishnamoorthy, S., 2014. Nutritional Quality in Freshwater Mussels, *Parreysia* spp. of Periyar River,Kerala, India; Research Journal of Recent Sciences, Vol. 3(ISC-2013), pp. 267-270.

- Shaikh, M. J., 2011. Seasonal variations in biochemical constituents in different body tissues of freshwater bivalve mollusc, lamellidens marginalis (lamark) from pravara river in Maharashtra; An international quaterly jornal on life scienc, 6(2), pp 297-299.
- Shakila and Nambudiri, 2012. Advances in harvest and post harvest technology in fish; New India publishing agency, New Delhi (India), pp. 307-335 .
- Sharma, S.K., 2006. Water quality assessment of Behta River using Benthic macro invertebrates. Life science journal, 3 (4), pp. 68-74.
- Shewfelt, R.L., 1981. Fish muscle lipolysis- a review. Journal of Food Biochemistry, 5(2), pp 79-100.
- Shrivastava, V.K., 1956. Benthic organisms of freshwater fish tank. Currr. Sci. 25,pp. 158-159.
- Shrivastava, V.K.,1959. Studies of freshwater bottom fauna, Quantitative composition and variation of the available food supply of fishes. Proc. Nat. Acad. Sci. India 29: 207-216.
- Singh, T. Y., Krishnamoorthy, M. and Thippeswamy, S., 2012. Seasonal changes in the biochemical composition of wedge clam, *Donax scortum* from the Padukere beach, Karnataka, Recent Research in Science and Technology, 4(12), pp. 12-17.
- Srilatha, G., Chamundeeswari, K., Ramamoorthy, K., Sankar, G., and Varadharajan, D., 2013. Proximate, Amino Acid, Fatty Acid and Mineral Analysis of Clam, *Meretrix casta* (Chemnitz) from Cuddalore and Parangipettai Coast, South East Coast of India., J Mar Biol Oceanogr 2:2, pp 2-7.
- Subba Rao, N.V. and Mitra, S.C., 1982. Bioecology of two Malaniid Snails (*Mollusca: Gastropoda*) in a pond near Calcutta. Bull. Zool. Surv. India, 34 (1 and 2) : 21-34.

- Subba, B.R. and Ghosh. T.K. 2000. Some freshwater molluscs from eastern and central Nepal. *J.Bombay nat. His. Soc.* **97(3)**: 452-455.
- Subba, B.R. 2003. Molluscan checklist of Ghodaghodi Tal area, Kailali district. *Our Nature* 1:1-2.
- Subba, B.R., R. Surana & K.P. Limbu. 2004. Report of molluscs from Chimdi lake (Birju Tal), Sunsari district, Nepal. *Our Nature* **2**: 45-46.
- Subba, B.R. and Pandey M.R.. 2005. *An inventory of aquatic biodiversity of Upper Mustang,Nepal*. Report submitted to King Mahendra Trust for Nature Conservation, ACAP,Upper Mustang Biodiversity Project. 54 p.
- Tanveer,A and Khan,D,1989. Seasonal Variations in the Environmental Factors & SnailPopulations in Four Different Habitats Around Lahore ; Dept. of Zoology of the Punjab University, Lahore: Punjab Univ. J. Zoo,Vol. 4 , pp 31-.69
- Tanveer, A., 1992. Response to temperature & heat tolerance of some snails Punjab university. *Bangladesh j.zoo.*20(2),:257-268.
- Thakur D.K., Mitra S.C. & Mitra S., 1992. Land and Fresh Water Molluscs (Part-II). Pub Handbook of Freshwater Molluscs of India. Calcutta: Zoological Survey of India. pp 1-128
- Uriarte, I. Fari, A., Hernandez, J., Scha, C. , Sorgeloos, P., 2004. Reproductive conditioning of Chilean scallop (*Argopecten purpuratus*) and the Pacific oyster (*Crassostrea gigas*): effects of enriched diets, *Aquaculture* (Elsevier) 230, pp 349–357.
- US Food and Drug Administration, 1998, Bacteriologicsal Analytical Manuals, 8th edn., Revision A. AOAC International, Gaithersburg, MD.

- Van Cleave, H.J. and Lederer, L.G., 1932. Studies on the life cycle of the snail, *Viviparus contectoides*. Journal of Morphology 53, pp 499-522.
- Vanugopalan, V. and M.A.James ,1969. Utilization of trash fish II. Studies on preparation of fish soup mix. Fish Technol., 6, pp 148-152.
- Vareltzis, K., Koufidis, D., Gavriilidou, E, Papavergou, E. and Vasiliadou, S., 1997. Effectiveness of a natural Rosemary (*Rosmarinus officinalis*) extract on the stability of filleted and minced fish during frozen storage. European Food Research and Technology 205, pp 93-96.
- Vedpathak,, A. N., 1989. Reproductive endocrinology of some Lamellibranch molluscs with special reference to environmental stress. Ph. D. Thesis, Marathwada University, Aurangabad (M.S.), India. pp.1 – 280.
- Yola, I. A. and Timothy, O.,2012. Proximate composition and consumer acceptability of african mudfish *Clarias gariepinus* smoked with two energy sources, ayero Journal of Pure and Applied Sciences, 5(2): 115 – 118
- Young, V.R.,1994. Adult amino acid requirements: the case for a major revision in current recommendations". *J. Nutr.* 124 (8 Suppl): 1517S–1523S.
- Wak, S.M., Myung, S.K., Lee, Y.J. and Seo, H.G., 2012. Efficacy of Omega-3 Fatty Acid Supplements (Eicosapentaenoic Acid and Docosahexaenoic Acid) in the Secondary Prevention of Cardiovascular Disease: A Meta-analysis of Randomized, Double-blind, Placebo-Controlled Trials. Archives of Internal Medicine 172 (9): 686–94.doi:10.1001/archinternmed.2012.262.PMID 22493407.
- Walne, P.R., 1970. The seasonal variation of meat and glycogen content od seven populations of oysters *Ostra edulis* L And a review of the literature Fisheries Investigation Series, 11,pp. 26-35.

- Wafer, M.V., Vijayaraghawan, M., S. and Krishnakumari, L., 1976. Seasonal changes in the nutritive values of the Green Mussel *Mytilus viridis* (L.). Indian J. Marine Sci., 5, pp. 252-254
- Weir, D.G. and Scott, J.M., 1999. Cobalamins Physiology, Dietary Sources and Requirements. In: Sadler M.J., Strain J.J., Caballero B., eds. Encyclopedia of Human Nutrition, 1: 394-401.
- Willett, W.C., 2007. The role of dietary n-6 fatty acids in the prevention of cardiovascular disease".J Cardiovasc Med 8: Suppl 1:S, pp. 42–45. doi:10.2459/01.JCM.0000289275.72556.13.PMID 17876199.
- WHO/FAO/UNU,2007. Protein and amino acid requirements in human nutrition, pub.:WHO Press (Geneva,Switzerland), pp 110-159.
- Yola, I. A. and Timothy, O.,2012. Proximate composition and consumer acceptability of african mudfish *Clarias gariepinus* smoked with two energy sources, ayero Journal of Pure and Applied Sciences, 5(2): 115 – 118
- Zhelyazkova, T.S., Pavlov, D. and Delchev, G., 2006. Content of amino acids in grain of winter pea, treated with plant growth regu-lators. In: *Proceedings of 4th International Eco-Conference on Safe Food*, Novi Sad, September 20-23 2006, pp. 343–349.

Web source :

- (1) <http://indiabiodiversity.org/species/show/224418>
- (2) <http://www.vitalhealthzone.com/nutrition/food-values/shellfish.html#9>
- (3) http://en.wikipedia.org/wiki/Pantothenic_acid
- (4) <http://umm.edu/health/medical/altmed/supplement/vitamin-b5-pantothenic-acid>
- (5) http://jn.nutrition.org/cgi/reprint/124/8_Suppl/1517S.pdf.
- (6) <http://www.wdcusick.com/protein.html -->>

- (7) www.buzzle.com/articles/essential-amino-acids-list.html -->*
- (8) <https://www.sickkids.ca/images/Research/Physiology-and-experimental-medicine/20022-Chris%20T%20PEM%20PP%20June%202009.pdf>*
- (9) <http://www.molluscs.at/gastropoda/index.html>*

* Extra reference