
List of Tables

| Tables | Page No. |
|---|-----------|
| Table 3.1. GC-MS analysis of methylated PS-I isolated from <i>T. heimii</i> . | 51 |
| Table 3.2. GLC-MS analysis of periodate-oxidised, reduced, methylated PS-I from <i>T. heimii</i> | 53 |
| Table 3.3. The ^1H NMR and ^{13}C NMR chemical shifts of PS-I. | 56 |
| Table 3.4. ROESY data for the PS-I isolated from the mushroom <i>T. heimii</i> . | 58 |
| Table 3.5. The ^{13}C NMR chemical shifts of Smith-degraded glycerol-containing disaccharide of the mushroom <i>T. heimii</i> in D_2O at $30\text{ }^\circ\text{C}$. | 59 |
| Table 4.1. GLC-MS analysis of methylated PS-II isolated from mushroom <i>L. fusipes</i> . | 72 |
| Table 4.2. GLC-MS analysis of periodate-oxidised, reduced, methylated PS-II. | 73 |
| Table 4.3. The ^1H NMR ^a and ^{13}C NMR chemical shifts of PS-II. | 75 |
| Table 4.4. ROESY data for the polysaccharide isolated from the mushroom <i>L. fusipes</i> . | 79 |
| Table 4.5. The significant $^3\text{J}_{\text{H,C}}$ connectivities observed in an HMBC spectrum for the anomeric protons/carbons of the sugar residues of PS-II from <i>L. fusipes</i> . | 81 |
| Table 4.6. The ^{13}C NMR chemical shifts of Smith-degraded glycerol-containing material of the mushroom <i>L. fusipes</i> in D_2O at $30\text{ }^\circ\text{C}$. | 83 |