



ISSN No. 0972-8503



Indian Journal of Biological Sciences

VOLUME - 24

April, 2018

INDIAN JOURNAL OF BIOLOGICAL SCIENCES
(A Peer Reviewed Journal)

Vol. 24

ISSN 0972-8503

2018

CONTENTS

	<i>Pages</i>
Evaluation of Lichen Diversity on Lianas in Selected Sacred Groves of West Midnapore District, West Bengal Uday Kumar Sen	1-12
Metabolic Study of Cycle Rickshaw Pullers Before and After Exercise Chandan Kumar Pradhan, Angshuman De, Indranil Chakraborty and Sridhar Thakur	13-21
A Novel Antibacterial Peptide from Whey-Based Fermented Product of Lactobacillus Ingluviei: Valorization of Whey Waste Arpita Mandal, Tanmay Paul, Arpita Patra, Animesh Samanta, Keshab Ch. Mondal and Dilip Kumar Nandi	22-30
A Comparative Study on the Impact of Thermal Working Environmental Factors and Workload on Cardiac Response Indicators in Male Food Crop Cultivators of Two Ethnic Groups Ayan Chatterjee, Sandipan Chatterjee, Surjani Chatterjee, Neepa Banerjee, Shankarashis Mukherjee	31-44
Customized Helmets for Two-Wheeler Users: An Anthropometric Study Satabdi Bhattacharje, Ayan Chatterjee, Neepa Banerjee, Sandipan Chatterjee, Shankarashis Mukherjee	45-50
Laccase Production by Potent Fungal Isolate Trichoderma Sp PUF 2 Through Submerged Fermentation and its Application in Dye Decolorization Priyanka Ghosh, Arpan Das and Uma Ghosh	51-59
Conservation of Ethno-Medicinal Plants Through Sacred Groves of Bankura District, West Bengal Atanu Nayak, Achintya Dey and Ram Kumar Bhakat	60-68
A Note on Ethno-Medicinal Herbs of Gopiballavpur Block-1 of Jhargram District, West Bengal, India Soma Dutta Das and Ram Kumar Bhakat	69-75
Aquatic and Marshland Dye Yielding Plants and Their Ethno-Medicinal Uses From Jhargram District, West Bengal, India Pijush Kanti Das	76-86
Evaluation of Extent of Mismatch Between Classroom Furniture and Anthropometric Measures of Girls' Students of High Madrasah (School) Benzir Parvin and Prakash C. Dhara	87-98
Isolation and Characterization of Melanoidins Degrading Bacteria from Sugar-Mill Effluent Ananda Kumar Saha, Md. Rabbi Al Zehad, Moni Krishno Mohanta Md. Fazlul Haque, Roksana Ara Ruhi and SK. Md. Atiqur Rahman	99-114
Recovery Effects of Zinc-Oxide and Fructooligosaccharides Nano Conjugate on Nicotine-Induced Oxidative Stress on Cardiac Tissue in Adult Albino Rat Monalisha Paul, Satarupa Bhattacharjee, Sukhan Das and Sandip Kumar Sinha	115-122

RECOVERY EFFECTS OF ZINC-OXIDE AND FRUCTOOLIGOSACCHARIDES NANO CONJUGATE ON NICOTINE-INDUCED OXIDATIVE STRESS ON CARDIAC TISSUE IN ADULT ALBINO RAT

Monalisha Paul¹, Satarupa Bhattacharjee¹, Sukhan Das² and Sandip Kumar Sinha^{1*}

¹Department of Human Physiology with Community Health, Vidyasagar University, Paschim Medinipur, West Bengal, Pin-721102

²Department of Physics, Jadavpur University, Kolkata, West Bengal, Pin-700032

ABSTRACT ■ Nicotine is major toxic component of tobacco products which are capable of producing free radicals and causes oxidative stress. Zinc and Fructooligosaccharides (one type of soluble dietary fiber) both are good for cardiac health as a dietary supplement (which is digested by gut bacteria). Both have a good and significant role on cardiac health. Our main objectives of this study to prepared a new nano-conjugate and apply against nicotine induced cardiac toxicity for recovery. In this study we prepared a nano-conjugate compound with zincoxide and Fructooligosaccharides (ZnO-FNC) by chemical synthesis method and characterized by using SEM and FTIR analysis. We apply ZnO-FNC on nicotine induced albino rat and measured SOD, LPO, CAT level of heart tissue and serum LDH level. Results shows that ZnO-FNC help to reduce oxidative stress of heart tissue induced by nicotine and all the data were changed significantly ($P < 0.05$) compare to nicotine treated group and closed to the control group. ZnO-FNC has potential effects on oxidative damage and it is a new approach of nutritional therapy against cardiac stress.

Key words: Zinc-oxide, Fructooligosaccharides, Oxidative stress. Heart

INTRODUCTION

After smoking, 90% nicotine absorbed very quickly through lung, easily reach in blood and heart through diffusion process and increased heart toxicity. In a normal person (non-smoker) nicotine level in plasma is 5 to 7 ng/ml utilised for nicotinic acid, melanin synthesis and other various metabolic purposes (Russell and Feyerabend, 1975). After smoking plasma nicotine level is 10 to 50 ng/ml (Matta et al., 2007) and causes oxidative stress, which is the main culprit for atherogenesis and thrombogenesis. Nicotine

reduced catalase and superoxide dismutase enzyme activity and increased H_2O_2 level in body fluid and increased lipid peroxidase level which causes degradation of cell membrane lipid and release free radicals. Soluble dietary fibers (SDF) is a non-digestible carbohydrates induced to promote healthy bacterial growth, responsible to relief from oxidative stress. Fructooligosaccharides (FOS) is one type of SDF, more effective for cardiac health and alleviate from hyper lipidemia, cardiac fat deposition, and oxidative stress than other SDF (Haubert et al., 2015). Normally zinc

* Corresponding author : e-mail: sinha_sandip@mail.vidyasagar.ac.in

diminishes the percentage of free radicals in cellular physiology and it has antioxidant property which is able to protect the cell membrane from oxidative stress (Marreiro et al., 2017). Zinc oxide nano particle (ZnONP) also acts as a scavenger of oxidative stress (Jacob and Rajiv, 2019). Here we prepared a new nano-conjugate form using zinc oxide and Fructooligosaccharides (ZnO-FNC) by chemical synthesis process in the field of nanobiotechnology. The main aims of this study was to investigate the potentiality of ZnO-FNC against nicotine induced oxidative stress in rat heart tissue, specifically alteration of catalase (CAT), superoxide dismutase (SOD), malondialdehyde (MDA) and lactose dehydrogenase (LDH).

2. MATERIALS AND METHODS

2.1 Synthesis and characterization of ZnO-FNC:

In the first step 3.8% (w/v) Fructooligosaccharide (Sigma Aldrich Company) was added with 0.025 mol zinc acetate (Merck Pharmaceutical companies) aqueous solution and 0.1 mol NaOH (Merck Pharmaceutical companies) mixed under constant stirring drop by drop for maintain pH (11) balance and keep it at room temperature for overnight (24 hour). Then the solution centrifuged at 2,000 rpm for 10 minutes. For removed NaOH, the supernatant water was discarded and washed the pallets three times by distilled water. Pallets were dried at 80°C then collect ZnO-FNC (Moharekar et al., 2014). ZnO-FNC sample was prepared in Department of Human Physiology with Community Health, Vidyasagar University, Paschim Medinipur, W.B (India). ZnO-FNC sample characterized through the FTIR (Fourier transforms infrared) spectra on Perkin Elmer Ltd software version 10.3.7 explain the chemical bonds are present in this conjugate sample. For FTIR

measurement we prepared pellet of ZnO-FNC with KBr 1 wt % and take spectrum (Vidyasagar University). The surface morphology and elemental analysis of the synthesized ZnO-FNC was done by field emission scanning electron microscopy (FESEM, FEI Inspect F50, and The Netherlands) Sigma model Carlzeics (Jadavpur University, West Bengal, India). The films of the sample were prepared on a carbon coated copy grid by just dropping a very small amount of the sample on the SEM grid which was then allowed to dry by putting it under a mercury lamp for 5 minutes (Weiss and Moser, 2015). From FESEM we understand the average practical size of ZnO-FNC.

2.2 Preparation of Nicotine and other applied sample solution: Nicotine (Purchase from Sigma Aldrich Company) was dissolved in 0.9% physiological solution (saline) and injected through intra-peritoneal (ip) according to effective dose of 3.5 mg/kg body weight one time daily 24 hour duration (Chattopadhyay and Chattopadhyay, 2008). On the basis of Tao et al., (2011) we prepared three doses of ZnO-FNC like 225 mg, 112.5 mg and 56.25 mg per kg body weight. In between them 112.5 mg / kg body dose of ZnO-FNC is more effective compare to other doses. Zinc oxide nanoparticle (ZnONP) and FOS (Fructooligosaccharids) both are purchase from Sigma Aldrich Company and both are apply as a same dose like 112.5mg/kg body weight. All the prepared samples like ZnONP, Fructooligosaccharids and ZnO-FNC are diluted in deionised water according to dose (112.5 mg/kg body weight) and stored at 2°C to 8°C.

2.3 Animal Care, grouping and mode of treatment:

Experiment was carried on male Albino rats (100-120 gm body weight) were obtained from were procured from SAHA ENTERPRIZE, 386 / 2, Nilachal, Birati, Kolkatta-700051, Email: sanjoy.animal2015@gmail.com. Regn.

No. 1828 / PO / BT / S / 15 / CPCSEA and keep them CPCSEA approved animal house of Vidyasagar University (Registration No. 2013/GO/Re/S/18/CPCSEA/2018) and ethical clearance from animal ethical committee (Ref. no. VU/IAEC/2/4). The animals were housed in a clean polypropylene cage in a standard temperature ($20 \pm 2^\circ\text{C}$) with relative humidity (45–60%) under 12-h light and dark cycles during the whole study period. Then animal were acclimatised in animal house for 15 days. Then animals were divided into 6 groups and each group having 10 animals. Throughout the experiments animals are provided normal standard balanced diet with water *ad-libitum*.

- **Group-I:** Control group : Animals were treated with 0.9% normal saline (i.p) for 14 days and next 21 days treated with distilled water (orally administrated) then scarified.
- **Group-II:** (Nicotine treated group) Animal were treated with nicotine (3.5 mg/kg BW) for 14 days (i.p) then scarified.
- **Group-III:** Auto-recovery group : This group were treated 14 days by nicotine (i.p) and next 21 days treated with distill water (orally administrated) then scarified.
- **Group-IV:** Zinc oxide nano particle (ZnONP) treated group : nicotine treated (i.p) for 14 days, next 21 days applied ZnONP (orally administrated) then scarified.
- **Group-V:** Fructooligosaccharides applied group : First 14 days treated with nicotine (i.p), next 21 days applied Fructooligosaccharides as a supplement (orally administrated) then scarified.
- **Group-VI:** ZnO-Fructooligosaccharides nano-conjugate (Zn-FNC)

applied group : First 14 days treated with nicotine (i.p), next 21 days applied Zn-FNC as a supplement (orally administrated) then scarified.

2.7 Tissue and serum collection: After scarifying, blood sample was collected by cardiac puncture technique in EDTA container and heart tissue stored in ice container. Here we use polytron homogenizer instrument (at 40°C) for tissue homogenate. 0.1gm heart tissue was homogenate in 1 ml phosphate buffer (0.05M, pH7) solution. The homogenate was centrifuged at 10,000 rpm for 20 min. The supernatant was used for the estimation of superoxide dismutase (SOD), catalase (CAT) and Lipid peroxidation (MDA) level on the other hand anticoagulant mixed blood sample was centrifuged at 3000 rpm for 10 minutes for separation of serum and it was utilized for the analysis of lactose dehydrogenase (LDH) activity.

2.8 Biochemical estimation:

2.8.1 Superoxide dismutase (SOD) estimation: 1mM diethylenetriaminepenta acetic acid added with 3ml Tris buffer (50mM, pH 8.2), 45 μl pyrogallol mixture (10mM pyrogallol added in 10mM HCl) and 10 μl of tissue supernatant mixed with aliquot of assay mixture and noted optical density (OD) at 420 nm. SOD activity was expressed as units/milligram protein (Marklund and Marklund, 1974).

2.8.2 Catalase (CAT) estimation: After homogenisation 0.1ml supernatant of tissue homogenate was added with 2.9 ml phosphate buffer (0.5M, pH 7) and then this mixture was transferred in a cuvette used as a control sample. In a cuvette 0.1ml supernatant of tissue homogenate also added with 1.9 ml phosphate buffer. Then mixed 1ml of H_2O_2 (11nm) solution with the previous aliquot assay mixture and immediately

observed the OD value at 240 nm with a one minute interval for 3 times (Beer and Sizer, 1952). Activity of catalase was calculated using following formula $\Delta A \times 10^3 / 40 \text{mg}$ protein in sample (U/mg protein).

2.8.3 Lipid Peroxidation (MDA) estimation: First of all 2 ml 0.375 % and 15% thiobarbituric acid and trichloroacetic acid were added respectively. Then 200 μl tissue supernatant and 1.0 ml distilled water were added with this mixture then boiled on a water bath at 95 °C for 20 min. After that the solution get pink colour and cooled under tap water. Then mixed 3 ml n-butanol to stop the internal reaction and observed OD value at 532 nm (Wills, 1969).

2.8.4 Lactose dehydrogenase (LDH) estimation: Serum LDH activity was estimated using Kit (Coral Clinical System, A Division of Tulip Diagnostic Pvt. Ltd., Vema, Goa, India). The LDH activity was expressed as Unit/L.

2.8.5 Protein estimation: The protein was determined by Lowry *et al.*, (1951) using cardiac tissue for the indirect measurement of enzyme.

2.9 Statistical analysis: All data are expressed as the Mean \pm Standard Error of Mean (SEM) and One Way Anova is used for statistical analysis by Origin 6.0 professional software (details shows in table).

3. RESULT AND DISCUSSION

3.1 Characteristics of ZnO-FNP:

In general, formation of NPs and their characterization were observed by following use of instruments. In this study, Field Emission Scanning Electron Microscopy (FE-SEM) analysis was done by using sigma model Carlzeics. SEM analysis is to provide supplied particle shape and size (nm). FE-SEM images of ZnO-FNC explain the structure is uniform spherical shaped and average particle size 40 nm which have shown in figure-1. The FTIR spectroscopy is also carried out to find out the purity and nature of NPs and bio-functionalized NPs synthesized by eco-friendly chemical synthesis method. In addition, it has been also showed the chemical bonds which were present in supplied sample. 493.19 denote the Zn-O stretching. 2101.45 gave CH_2 and 2346.58 give the CH_3 stretching. 3389.76 to 3786.54 indicate the O-H stretching (Silverstein *et al.*, 1981) shows in figure 2.

3.2 Effects of ZnO-FNC on oxidative stress:

Nicotine, the major component of tobacco product plays an important role in the development of lung, heart and liver complications. It causes oxidative damage in these organs. As we know that nicotine it is a potential oxidant, which is capable of producing free radicals and reactive oxygen

Table 1. Shows the antioxidant and oxidative stress markers in different groups (Mean \pm SEM). a, b, c d, e and f different letters denote significance within control groups or treatment groups (enteritis) at $P < 0.05$. (One way ANOVA)

	Group I	Group II	Group III	Group IV	Group V	Group VI
SOD (U/mg protein)	4.32 ^a \pm 0.37	1.56 ^{bce} \pm 0.26	1.46 ^{bce} \pm 0.26	1.97 ^{ce} \pm 0.29	1.68 ^d \pm 0.26	3.92 ^a \pm 0.56
MDA (nmol/ gm tissue)	12.84 ^a \pm 1.81	15.97 ^{bc} \pm 1.84	15.60 ^{bc} \pm 1.25	14.72 ^b \pm 1.98	15.63 ^c \pm 1.30	13.29 ^a \pm 1.93
CAT (U/mg protein)	42.20 ^a \pm 1.45	19.44 ^{bc} \pm 1.66	22.37 ^{bdc} \pm 1.43	24.68 ^{dc} \pm 1.64	21.84 ^c \pm 1.44	35.17 ^e \pm 2.61
LDH(U/L)	436.27 ^a \pm 2.82	1130.86 ^b \pm 3	913.54 ^c \pm 2.69	486.57 ^d \pm 2.5	673.42 ^e \pm 3.2	365.81 ^f \pm 2.60




Fig. 1: Field Emission Scanning Electron Microscope image of synthesized Zn-FNC




Fig. 2: Fourier transform infrared spectrum of Zn-FNP.

species. So inside the cell, nicotine-induced free radicals react with bio-membranes causing oxidative destruction of polyunsaturated fatty acids and forming cytotoxic (Chiagoziem et al., 2014) components. An in-vitro cytotoxicity study on malaria parasites suggested that infected HeLa cell shows the application of chitosan-tripolyphosphate (CS-TPP) nanoparticles (NPs) conjugated with chloroquine help to reduced oxidative stress (SOD, CAT, LDH and MDA level). Chitosan is a type of dietary fiber

(Tripathy et al., 2012) responsible for controlling various types of physiological functions. CAT is an antioxidant as well as dismutase enzyme found in living aerobic organism such as peroxisomes and mitochondria, specially splitting hydrogen peroxide into water and oxygen. On the other hand SOD transfer the electron from one molecule of the superoxide anion to another and form hydrogen peroxide from free radical $2O_2^{\cdot-} + 2H^+ \rightarrow H_2O_2 + O_2$. (Ma et al., 2012). Table-1 shows the CAT and SOD level of

nicotine treated group which are significantly ($p < 0.05$) decrease than the control groups. In an auto-recovery phase body able to recover the oxidative stress but very slowly. Whereas ZnO-FNC shows very quick recovery response for the level of CAT and SOD in heart tissue ($p < 0.05$) which are approximately similar to control groups. Various findings reported that specially free radicals (allylic hydrogen atom) initiated lipid peroxidation (LPO) and breakdown lipids of cell membranes (Snezana et al., 2014). Lipid peroxidation has been introduced in pathogenesis of a number of diseases (Chiagoziem et al., 2014). Nicotine increased lipid peroxidation and MDA production (MDA an end product of lipid peroxidation). Wide range of MDA production may causes ather osclerosis, stroke, diabetes, and aging (Landau et al., 2013). After 15 days of nicotine treatment, MDA level was 15.974 ± 1.8416 (nmol/ gm tissue, $p < 0.05$) whereas in control groups 12.48 ± 1.8148 (nmol/ gm tissue, $p < 0.05$). In case of ZnO-FNC group (group-VI) after the end 20 days of treatment, MDA level of heart tissue reduce significantly ($p < 0.05$) at 13.29 ± 1.93497 (nmol/ gm tissue). Form these result it is very clear that ZnO-FNC skilfully reduce the free radicals and it may have some antioxidant activity. Lactate dehydrogenase (LDH) is a another type of glycolytic enzyme found in various tissues like skeletal muscle, heart, liver, kidneys, brain, lungs, and red blood cells. The intensity of oxidative stress induces LDH release from the tissues and raises its level in the blood causes of stroke, cancer, heart attack, blood flow deficiency, haemolytic anaemia, hepatitis, muscle injury and tissue death (Berridge et al., 2013). Iso-enzymes of LDH like LDH-1 & LDH-2 are responsible for myocardial infection and LDH-5 responsible for liver disease (Usher et al., 1974). For nicotine toxicity LDH-1 type is exposed

(Gamieldien et al., 2008). At the same time both zinc and SDF are responsible for the development of intestinal lactic acid and acetic acid producing bacterial growths which are capable to reduced LDH level (Aldunate et al., 2015 and Leonardi et al., 2013). Hamza et al., (2018) pointed out that green tea leaves extract and zinc oxide nano practical complex form potentially reduced the LDH level of liver tissue. In case of our study we found that nano-conjugate supplement (ZnOFNC) is capable for recovery of oxidative damage by lowering LDH and MDA level in 20 days (group-VI) treatment towards the control group. Therefore we may be suggested that nano conjugate dietary supplement (ZnOFNC) have a significant beneficial effect on nicotine induced oxidative damage in biological system, which may be important for the prevention of smoking (nicotine) induced cardiovascular damage for smokers.

4. CONCLUSION:

So in this present studies we try to assess the toxic effect of nicotine on cardiac tissue through the measurement of CAT, SOD, LPO and LDH level. Zinc oxide and fructooligosaccharides nano-conjugate (ZnOFNC) efficiently diminished the oxidative stress, free radical scavenging activity and antioxidant properties within few days. We are also trying to investigate the other biomarkers related to nicotine induced cardiovascular damage in future.

5. ACKNOWLEDGEMENT:

This study was supported by Vidyasagar University, Midnapore for providing the financial support through personal research grant (PRG) and Department of Physics, Jadavpur University West Bengal, India for their investigation support through instrumentation.

6. REFERENCES

- Aldunate, M., Srbinovski, D., Hearps, A.C., Latham, C.F., Ramsland, P.A., Gugasyan, R., Cone R.A and Tachedjian G. (2015): Antimicrobial and immune modulatory effects of lactic acid and short chain fatty acids produced by vaginal microbiota associated, *Front Physiol.* 2 (6):164.
- Beers, R.F.Jr. and Sizer, I.W. (1952): A spectrophotometric method for measuring the breakdown of hydrogen peroxide by catalase, *J Biol Chem.* 195(1):133-40
- Berridge, R. B., John F, Vleet, V. and Herman, E. (2013): Cardiac, Vascular, and Skeletal Muscle System, Haschek and Rousseaux's Handbook of Toxicologic Pathology. 3: 1567-1665.
- Chattopadhyay, K. & Chattopadhyay, B.D. (2008): Effect of nicotine on lipid profile, peroxidation & antioxidant enzymes in female rats with restricted dietary protein, *Indian J Med Res.* 127: 571-576 of maternal nicotine exposure during gestation and lactation on lactate dehydrogenase
- Chiagoziem, A. O., Sunny, O. A., Victoria, I. O., Azeezat, L. A. and Oluwafemi, E. K. (2014): Protective Effect of Curcumin against the Liver Toxicity Caused by Propanil in Rats. Pbl. Hindawi Publishing Corporation, International Scholarly Research Notices. Article ID 853697, 8 pages.
- Gamielien, K. and Maritz, G. (2008): Influence isoenzyme profile and transcript levels in developing neonatal rat lung, *Pathophysiology.* 15(1):1-8.
- Hamza Z.R., Salmi A.A.F. and Shenawy S.E.N. (2018): Nanoparticles Effects on Zinc Oxide/green Tea Complex on the Lipid Profile and Liver Functions of Rats after Monosodium Glutamate Treatment. *Journal of Applied Sciences* 18(2):65-70.
- Haubert, N.J.B.G.B., Marchini J. S., Cunha S.F.C. and Suen V.M.M. (2015): Choline and Fructooligosaccharide: Non-alcoholic Fatty Liver Disease, Cardiac Fat Deposition, and Oxidative Stress Markers, *Nutrition and Metabolic Insights.* 4(8):1-6.
- Jacob, V. and Rajiv, P. (2019): In vitro analysis: the antimicrobial and antioxidant activity of zinc oxide nanoparticles from curcuma longa, *Asian, J Pharm Clin Res.* 12(1): 200-204.
- Landau, G, Kodali, K.V, Dr Malhotra, J. D., and Kaufman, R. J.K. (2013): Detection of Oxidative Damage in Response to Protein Misfolding in the Endoplasmic Reticulum, *Methods in Enzymology.* 526 : 231-250.
- Leonardi, A., Zanoni, S., Lucia, D. M., Amaretti, A., Raimondi, S., and Rossi, M. (2013): Zinc Uptake by Lactic Acid Bacteria. *ISRN Biotechnology.* Article ID 312917, 5 pages.
- Lowry O H., Rosebrough J N., Farr A. L, and Randall R J. (1951): Protein measurement with the folinphenol reagent. *J. Biol. Chem* 193: 265
- Ma, Q., He, X. (2012): Molecular basis of electrophilic and oxidative defense: promises and perils of Nrf2, *Pharmacol Rev.* 64:1055-81.
- Marklund, S., Marklund, G. (1974): Involvement of superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase, *Eur. J. Biochem.* 47: 469-474.
- Marreiro, D.D., Cruz, K.J., Morais, J.B., Beserra, J.B., Severo, J.S., and de Oliveira, A.R. (2017): Zinc and Oxidative Stress: Current Mechanisms, *Antioxidants (Basel).* 29; 6(2).
- Moharekar S., Raskar P., Wani A. and Moharekar S. (2014): Synthesis and comparative study of zinc oxide nanoparticles with and without capping of pectin and its application. *World Journal of Pharmacy and Pharmaceutical Sciences.* 3 (7):1255.
- Russell, M.A., and Feyerabend, C. (1975): Blood and Urinary nicotine in non-smokers, *Lancet.* 1(7900):179-81.
- Matta S.G., Balfour D.J., Benowitz N.L., Boyd R.T., Buccafusco J.J., Caggiula A.R., Craig C.R., Collins A.C., Damaj M.I., Donny E.C., Gardiner P.S., Grady S.R., Heberlein U., Leonard S.S., Levin E.D., Lukas R.J., Markou A., Marks M.J., McCallum S.E., Parameswaran N., Perkins K.A., Picciotto M.R., Quik M., Rose J.E., Rothenfluh A., Schafer W.R., Stolerman I.P., Tyndale R.F., Wehner J. Mand Zirger J.M. (2007): Guidelines on nicotine dose selection for in vivo research, *Psychopharmacology.* 190:269-319.
- Silverstein, R.M., Bassler, G.C., and Morrill, T.C. (1981): *Spectrometric Identification of Organic Compounds*, 4th ed. New York: John Wiley and Sons. QD272.S6 S55.

- Snezana, A. J. and Bojan, Z. S. (2014): Cadmium and lipid peroxidation, *Vitamins & Hormones*. 4C:391-425.
- Tripathy S., Das S., Chakraborty P. S., Sahu K. S., Pramanik P. and Roy S. (2012): Synthesis, characterization of chitosan–tripolyphosphate conjugated chloroquine nanoparticle and its in vivo anti-malarial efficacy against rodent parasite: A dose and duration dependent approach, *International Journal of Pharmaceutics*. 434 (2012): 292– 305
- Usher, D.J., Shepherd, R.J. and Deegan, T. (1974): Serum lactate dehydrogenase isoenzyme activities in patients with asthma, *Thorax*. 29(6):685-9.
- Weiss AC and Moser D R. (2015): Sample Preparation of Nano-sized Inorganic Materials for Scanning Electron Microscopy or Transmission Electron Microscopy. Scientific Operating Procedure SOP-P-2, Environmental Consequences of Nanotechnologies. ERDC/GSL SR-15-1. 8-10.
- Wills, E.D. (1969): Lipid peroxide formation in microsomes. Relation of hydroxylation to lipid peroxide formation, *Biochem. J.* 113: 333–341.
- Yi Tao, Hongliang Zhang, Bing Gao, Jiao Guo, Yinming Hu, and Zhengquan Su. (2011): Water-Soluble Chitosan Nanoparticles Inhibit Hypercholesterolemia Induced by Feeding a High-Fat Diet in Male Sprague-Dawley Rats, *Journal of Nanomaterials*. 814606: 5.

INDIAN JOURNAL OF BIOLOGICAL SCIENCES

INSTRUCTION TO AUTHORS

Original and unpublished manuscripts should be submitted in triplicate, typed double spaced on one side of the paper (A4 bond) with 3cms, margin on all sides. Along with the hard copy the paper is to be submitted in a CD. **The paper may preferably be submitted electronically by the following email: ijbsvu@gmail.com.**

Full length research paper:

The arrangement of the manuscript should be as follows: Title Page, Abstract, Key words, Introduction, Methods, Results, Discussion, Acknowledgments, References, Tables, Figures, Legends of Figures, Full length of paper should not exceed 10 printed pages.

Title page: It should contain the following information:

- (i) The title of the paper which should be concise but informative, (ii) a short running head of not more than 10 words placed at the foot of title page, (iii) first name, middle initial and last name of each author,
- (iv) Name of department(s) and institution(s) which the work should be attributed, (v) name, address and email of author for correspondence.

Abstract: The second page should carry an abstract of not more than 200 words. The abstract should state the purpose of the study, basic procedure, main findings and principal conclusions. Abstract should be followed by relevant key words (not more than five)

Introduction: This should contain a concise statement of the purpose of the article. Only pertinent references should be given. Reviewing of the literature should not surpass what is essential to indicate the motive for the research undertaken and the important background.

Methods: The methodology, apparatus and procedure in sufficient detail should be identified to allow other workers to repeat the experiments. Standard methods can, however, be identified by proper references. The new or substantially modified methods should be described giving reasons for using them.

Results: This section should be clear and concise, and may be combined with the Discussion section if appropriate. The Results section should describe only the most important results of the study and include as vital discussion as possible. Measures of statistical significance must be clearly indicated, and tables and figures should be cited in numerical order.

Discussion: This should precisely deal with interpretation of results. Emphasis should be given on the new and important aspects of the study and conclusions that follow from them. Recommendations, when appropriate, may be included.

References: In the text reference should be cited with author's surname and year of publication in parentheses. The full reference should be cited alphabetically according to the surnames of the authors at the end of the manuscript. The following forms of citations should be used.

Journals: Bunt, A.H., Lund, R.H and Lund, J.S. (1994): Retrograde axonal transport of the albino rat Retina, Brain. Res. 72: 215-228.

Books: Campbell, A.M. (1984): Monoclonal Antibody Technology. ELSEVIER, Amsterdam, pp 5066.

Individual Chapters in book: Bladen, G.E. (1975): Absorption of fluid and electrolytes in health and disease. In: Intestinal Absorption in Man (I. McColl and G.E. Bladen, eds.), Academic Press. London, 135-146. Journal title should be abbreviated as in Index Medicus or Biological Abstract.

Tables: Tables should be typed on separate sheets using double space. Each table should be numbered (Roman numerals) consecutively with a brief caption on the top of the table.

Illustration: Figure should be numbered and marked on the back. Line drawing sufficiently thick for good reproduction should be made with Chinese ink on plane white paper or tracing paper.

Legends of the figures should be type written on separate sheets and their positions in the text should be indicated in the manuscript.

Computerized illustrations should have good resolution (at least 600 pixels x 600 pixels at 300 dpi). Photographs should usually be in black and white. However, colour photograph may be accepted, if it is necessary for illustration. A nominal amount of money may be charged from the authors for printing colour photograph or figures. A clear print in glossy paper, large enough to be legible after 25% reduction, is necessary for reproduction. These should be submitted along with the paper.

Short Research Communication presents a brief study, or sometimes prelude but original research findings that might be less extensive than a full research paper. Short Research Communication is limited to 2000 words. It should have a set of keywords and an abstract summarizing background of the work, the results and their implications. Results and Discussion section should be combined and followed by Conclusion. Materials and Methods will remain as an individual section. The number of references is limited to 30 and the number of figures and/or tables combined is limited to 4.

Review or mini-review should be convincing and of high interest. A minimum of two figures/illustrations should be included in the review or mini-review that should be some 3000 or 4000 words long (excluding references and figure legends).

All official correspondence should be made in the following address: The Editor-in-chief, Indian Journal of Biological Science, Department of Human Physiology with Community Health, Vidyasagar University, Midnapore-721102, West Bengal or preferably by email (ijbsvu@gmail.com)

INDIAN JOURNAL OF BIOLOGICAL SCIENCES

(A Peer Reviewed Journal)

a publication of Vidyasagar University, which accepts original unpublished articles, review papers and short communications in the field of biological sciences.

Editorial Board

Editor-in-Chief : Prakash C. Dhara

Editor : Ram Kumar Bhakat, Jayanta Kumar Kundu, Kaushik Bose

Advisory Committee

Internal Members

1. S. K. Chakraborty
2. B.C. Patra
3. D. Ghosh
4. F. Chakraborty

External Members

1. N. C. Dutta
2. N. D. Paria
3. D. J. Chatterjee
4. A. M. Chandra
5. A. Basu
6. R. K. Das
7. T. K. Ghosh
8. A. Bhattacharjee

**Direct Manuscripts, Comments on Articles / Papers and
Correspondence to**

Professor Prakash C. Dhara

Editor-in-Chief

Indian Journal of Biological Sciences

Department of Human Physiology with Community Health

Vidyasagar University

Midnapore - 721 102

West Bengal

India

email : ijbsvu@gmail.com

The concerned authors are responsible for the views expressed, findings made and conclusions drawn in the articles.

Price : Rs. 50/-

Published by the Registrar, Vidyasagar University, Medinipur - 721 102, West Bengal, India
and Printed by Giri Printers, Mob- 9434005458, Midnapore - 721 102.