

## *List of Tables*

<i>Table</i>	<i>Title</i>	<i>Page</i>
<i>Chapter III</i>		
3.1	Fluorescence lifetime of $P^1$ in THF and its aggregated hydrosols (sample-c) of $P^1$ . (excitation:336 nm)	47
3.2	Electrophilic $f^+$ and nucleophilic $f^-$ condensed Fukui functions and second order Fukui parameter, $f^{(2)}(r)$ of Benz(a)anthracenemolecule calculated using DFT B3LYP 6-31G (d, p) level of theory.	49
3.3	Computed value of $f^{(2)}(r)$ of possible interacting atomic centers in Benz(a)anthracene dimer.	50
<i>Chapter V</i>		
5.1	Stern-Volmer quenching constant ( $K_{SV} M^{-1}$ ) for different nitroaromatics.	84
<i>Chapter VI</i>		
6.1	Stern-Volmer quenching constant ( $K_{SV} M^{-1}$ ) for different nitroaromatics.	108
6.2	A comparative study of our work with the other reported works in the literature with reference to $K_{SV}$ , Detection limit and Anti interference study.	108
<i>Chapter VII</i>		
7.1	Photophysical parameters of $P^5$ , Al- $P^5$ , Zn- $P^5$ and aggregated $P^5$ .	122
7.2	Comparative Study	135-136

## *List of Schemes*

<i>Scheme</i>	<i>Title</i>	<i>Page</i>
<i>Chapter I</i>		
1.1	A) Schematic representation of the structural description of	8

	Herringbone as well as H- and J-aggregates in terms of relative translation of molecules. B) Exciton splitting in J-aggregate and H-aggregate.	
1.2	Schematic representation showing binding of analyte (guest) by a chemosensor (host), producing a complex accompanying with change in optical properties.	12
1.3	Design of target molecules	16
<i>Chapter II</i>		
2.1.1-- 2.1.8	Schematic representation of various types of AIE/AIEE based Low dimensional material	18-22
2.2.1-- 2.2.9	Schematic representation of various types of fluorescent chemosensors for 2,4,6-Trinitrophenol.	23-30
2.3.1-- 2.3.8	Schematic representation of various types of fluorescent Dual chemosensors for Al (III) and Zn (II) Ions.	31-36
<i>Chapter III</i>		
3.1	Schematic representation of P <sup>1</sup> with its long (a, a') axis and short (b, b') axis.	44
3.2	Stacked conformer of P <sup>1</sup> dimer.	50
<i>Chapter IV</i>		
4.1	Schematic representation of P <sup>2</sup> with its long (a, a') and short (b, b') axis.	65
<i>Chapter V</i>		
5.1	Synthesis of 4-[(2-Hydroxy-naphthalen-1-ylmethylene)-amino]-1,5-dimethyl-2-phenyl-1,2-dihydro-pyrazol-3-one (P <sup>3</sup> )	70
<i>Chapter VI</i>		
6.1	Proposed Mechanistic Pathway for Aggregation Induced Emission Enhancement Behavior of (P <sup>4</sup> ).	97
6.2	The proposed mechanism of P <sup>4</sup> for sensing H <sup>+</sup>	98
<i>Chapter VII</i>		
7.1	Synthesis of 4-[(2-Hydroxy-3-methoxy-benzylidene)-amino]-1,5-dimethyl-2-phenyl-1,2-dihydro-pyrazol-3-one (P <sup>5</sup> )	113