2019

B.Sc. (Honours)

5th Semester Examination

PHYSICS

Paper - C11T

Full Marks: 40

Time: 2 Hours

The figures in the margin indicate full marks. Candidates are required to give their answers in their own words as far as practicable.

- 1. Answer any *five* questions from the following: $5\times2=10$
 - (a) Using relation $[\hat{x}, \hat{p}] = i\hbar$, prove

$$\left[\hat{p},f(x)\right]=-i\hbar\frac{\partial f}{\partial x}.$$

(b) The eigenvalue of an operator $\left(x + \frac{d}{dx}\right)$ is α . Find eigenfunction.

[Turn Over]

- (c) An electron has a speed of 300 m/s, accurate to 0.01%. With what accuracy can the position of the electron be located?
- (d) Write down the Hamiltonian of the electron in the hydrogen atom.
- (e) In a many electron atom, the orbital, spin and total angular momenta are denoted by \overline{L} , \overline{S} and \overline{J} respectively. If L=2, S=1 and J=2, find the angle between \overline{L} and \overline{S} using vector atom model.
- (f) A spectral line of wavelength 4500Å when produced in a magnetic field to 10 Tesla is observed to split to a normal zeeman triplet. Calculate the wavelength separation between components of the triplet.
- (g) Compute the value of $\langle r \rangle$ in the ground state of hydrogen atom.
- (h) Find Landé-g-factor and total magnetic moment for ${}^6S_{5/2}$ given $\mu_B = 9.27 \times 10^{-24} \ J/T$

- 2. Answer any *four* questions from the following: $5\times4=20$
 - (a) What is the physical significance of normalization of wave function? Find normalization const. A and sketch the graph of the function.

$$\psi(x) = A(x-1)x, \quad 0 \le x \le l.$$
 1+2+2

- (b) (i) What is meant by space quantisation?
 - (ii) In a Stem-Gerlach experiment, the gradient of magnetic field is 5 volt. s.m⁻²/mm with pole pieces of 0.07m long. A narrow beam of silver atoms from an over at 1000K passes through the magnetic field. Calculate the separation of the beams as they emerge from the magnetic field.
- (c) (i) What do you mean by stationary states?
 - (ii) The wave function of a particle in a stationary state with energy E_0 at time t = 0 is $\psi(x)$. After now much minimum

[Turn Over]

time will the wave function be again $\psi(x)$? 2+3

(d) A free particle which is initially localized in the range -a < x < a is released at time t = 0.

$$\psi(x,0) = \begin{cases} A & \text{if } -a < x < a \\ 0 & \text{otherwise} \end{cases}$$

Calculate $\phi(x)$ {wave-vector function of $\psi(x, 0)$ in momentum space} and find $\psi(x, t)$.

(e) Write down the Hamiltonian and Schrödinger wave equation for hydrogen atom. Separate the Schrödinger equation for the motion of electron in hydrogen atom into radial and angular parts.

(f) Discuss briefly, with theory, the Stren-Gerlach experiment.

Justify the use of a beam of silver atoms in the experiment.

3. Answer any one question from the following:

 $10 \times 1 = 10$

- (a) (i) Describe the theory of anomalous Zeeman effect.
 - (ii) Illustrate with appropriate diagrams the Zeeman splitting of sodium D_1 and D_2 lines.
 - (iii) Calculate the processional frequency of an electron orbit when placed in a magnetic field of strength 4T. 5+3+2
- (b) (i) The potential energy of a linear harmonic oscillator consisting of a particle of mass m and executing oscillations with an angular frequency ω_0 is $V(x) = \frac{1}{2}m\omega_0^2x^2$.

Write down Hamiltonian operator and time independent Schrödinger equation for the oscillator. What are the eigenvalues and eigenfunctions of the Hamiltonian? 2+3

- (ii) What is Zero-point energy? How does it reconcile with classical view point? 2+2
- (iii) Is it possible for a free particle to have negative energy?