

2019

B.Sc.

1st Semester Examination

PHYSICS (Honours)

Paper—GE 1-P

Full Marks : 20

Time : 3 Hours

*The figures in the margin indicate full marks.
Candidates are required to give their answers
in their own words as far as practicable.*

Elements of Modern Physics Lab

1. Determine the value of Boltzmann constant using V-I characteristics of p-n diode [value of n is to be supplied]
 - (a) Working formula. 2
 - (b) Circuit diagram. 2
 - (c) Implementation of the circuit. 2
 - (d) Data for $\ln I_n$ Vs. V curve. 5
 - (e) Drawing $\ln I_n$ Vs. V curve. 2
 - (f) Calculation of k from graph. 2
2. Determine work function of the material filament of directly heated vacuum diode. [Temperature

[Turn Over]

co-efficient of resistance of filament material is to be supplied]

- | | |
|--|---|
| (a) Working formula. | 2 |
| (b) Circuit diagram. | 2 |
| (c) Implementation of the circuit. | 2 |
| (d) Data for $\ln \frac{1}{T^2}$ Vs. $\frac{1}{T}$ graph. | 5 |
| (e) Drawing $\ln \frac{1}{T^2}$ Vs. $\frac{1}{T}$ graph. | 2 |
| (f) Calculation of work function from graph. | 2 |
| 3. Determine the wave length of H_{α} emission line of Hydrogen atom using grating method. [Number of rulings per mm is to be supplied] | |
| (a) Working formula. | 2 |
| (b) Vernier constant of the spectrometer. | 1 |
| (c) Setting the grating for normal incidence. | 2 |
| (d) Reading for deviation of H_{α} line for any order on either side of the central maximum. (Reading for both vernier are to be taken) | 8 |
| (e) Calculation of the wavelength of H_{α} emission line. | 2 |
| 4. Determine the wave lengths of absorption lines in the rotational spectrum of Iodine vapour. | |
| (a) Working formula. | 2 |

- (b) Vernier constant of the spectrometer. 1
- (c) Setting the grating for normal incidence. 2
- (d) Reading for deviation of H_{α} line for any two absorption lines specified by the examiner. (Readings on Either side of the central maximum and for one vernier is to be taken) 8
- (e) Calculation of the wavelength of H_{α} emission line. 2
5. Study the diffraction patterns of a single slit using laser and measure its intensity variation using LDR.
- (a) Theory 2
- (b) Data of LDR current versus angle of diffraction (only main scale reading with one vernier) on each sides of the central maximum. 8
- (c) Drawing intensity (in arbitrary unit) versus angle of diffraction graph. 3
- (d) Calculation of slit width from position of 1st order minima. 2
6. Determine of Planck's constant by photo electric effect.
- (a) Working formula. 2
- (b) Figure with circuit diagram. 3
- (c) Data for stopping potential of photo diode for lights of five wavelengths. 5

[Turn Over]

- (d) Drawing stopping potential versus wavelength graph. 3
- (e) Calculation of Planck's constant from graph. 2
7. Determine the value of $\frac{e}{m}$ by magnetic focusing (Thomson's method)
- (a) Working formula.
- (b) Calculation of magnetic field of the Helmholtz coil from magnetizing current. [Radius and number of turns of the Helmholtz coil are to be supplied] 2
- (c) Data for D^2 versus V graph (for at least four accelerating voltages) 7
- (d) Drawing D^2 versus V graph. 2
- (e) Calculation of $\frac{e}{m}$. 2

Experiment 15 marks

Laboratory Note book 02 marks

Viva voce 03 marks
