

List of figures

Figure 1.1: Typical structures of novolac and resole resin

Figure 1.2: Chemical structures of some typical azo-dye colorants

Figure 1.3: Reductive-cleavage of Direct Red 28 using an azo-reductase enzyme

Figure 1.4: Schematic diagram of nanoparticles (NPs) synthesis (top-down and bottom-up methods)

Figure 2.1 The structures of target novolac-based polymeric networks

Figure 2.2. Reaction mechanism of crosslinking of novolac epoxy resin with ethylenediamine.

Figure 2.3 ^1H NMR spectrum of **2.1**

Figure 2.4 ^1H NMR spectrum of **2.2**

Figure 2.5 ^{13}C NMR spectrum of **2.2**

Figure 2.6 FTIR spectra of (a) **2.3** (b) **2.4**

Figure 2.7 Solid state ^{13}C NMR spectra of (a) **2.3** and (b) **2.4**

Figure 2.8 Scanning electron micrograph of **2.3**

Figure 2.9 Scanning electron micrograph of **2.4**

Figure 2.10 Traces of thermogravimetric analysis (TGA / DTG) of (a) network polymer **2.3** and (b) network polymer **2.4** (Heating rate of $10^\circ\text{C}/\text{min}$ under nitrogen flow ($100\text{ ml}/\text{min}$)).

Figure 2.11 Swelling studies of polymer networks **2.3** and **2.4**.

Figure 2.12. Size distributions of the network powders **2.3** and **2.4**

Figure 2.13. UV-vis spectral changes of the solutions of (a) MO ($7.64 \times 10^{-5}\text{M}$), (b) OG ($2.21 \times 10^{-4}\text{M}$) and (c) OII ($3.06 \times 10^{-4}\text{M}$) as a function of adsorption time for **2.4**; (pH 7.20, different time intervals, 25°C).

Figure 2.14. The effect of adsorption time on MO adsorption capacity of **2.4**

Figure 2.15. The effect of adsorption time on OII adsorption capacity of **2.4**

Figure 2.16. Effect of pH on the azo dye adsorption capacity of **2.3** and **2.4** (a, d: pH = 2.30; b, e : pH = 7.20; c, f: pH = 10.96; t = 48h; T= 298 K); MO adsorption ($C_0 = 7.64 \times 10^{-5}M$).

Figure 2.17. Effect of pH on the azo dye adsorption capacity of **2.3** and **2.4** (a, d: pH = 2.30; b, e: pH = 7.20; c, f: pH = 10.96; t = 48h; T= 298 K); OII adsorption ($C_0 = 3.06 \times 10^{-4}M$).

Figure 2.18. Effect of pH on the azo dye adsorption capacity of **2.3** and **2.4** (a, d: pH = 2.30; b, e: pH = 7.20; c, f: pH = 10.96; t = 48h; T= 298 K); OG adsorption ($C_0 = 2.21 \times 10^{-4}M$).

Figure 2.19. Freundlich isotherms for the adsorption of MO onto **2.3** at (a) pH 2.30 (-●-) and (b) pH 7.20 (-▼-) at 25°C

Figure 2.20. Freundlich isotherms for the adsorption of MO onto **2.4** at (a) pH 2.30 and (b) pH 7.0 at 25°C

Figure 2.21. Freundlich isotherms for the adsorption of OII onto **2.3** at (a) pH 2.30 and (b) pH 7.20 at 25°C

Figure 2.22. Freundlich isotherms for the adsorption of OII onto **2.4** at (a) pH 2.30 and (b) pH 7.20 at 25°C

Figure 2.23. Freundlich isotherms for the adsorption of OG onto **2.3** at (a) pH 2.30 and (b) pH 7.0 at 25°C

Figure 2.24. Freundlich isotherms for the adsorption of OG onto **2.4** at (a) pH 2.30 and (b) pH 7.0 at 25°C

Figure 2.25 A proposed mechanism of adsorption: chemical interactions occurring within the network adsorbents

Figure 2.26 Azo dye (MO) adsorption for 48h onto network polymer matrix associated with visual color change

Figure 3.1. Iron(III) loaded novolac-based networks **3.1** and **3.2**

Figure 3.2 FTIR spectrum of **3.1**

Figure 3.3 FTIR spectrum of **3.2**

Figure 3.4 The XRD pattern of **3.2**

Figure 3.5 The effect of adsorption time on azo-dye adsorption of **3.1** and **3.2** at pH = 7.20 (a) MO adsorption ($C_0 = 50 \text{ mg L}^{-1}$; $1.52 \times 10^{-4}\text{M}$); (b) OG adsorption ($C_0 = 150 \text{ mg L}^{-1}$; $3.31 \times 10^{-4}\text{M}$).

Figure 3.6 Azo dye adsorption capacity of **3.1** and **3.2** (pH = 7.20; t = 48h; T= 298 K); (a) MO adsorption ($C_0 = 50 \text{ mg L}^{-1}$; $1.52 \times 10^{-4}\text{M}$); (b) OG adsorption ($C_0 = 150 \text{ mg L}^{-1}$; $3.31 \times 10^{-4}\text{M}$)

Figure 3.7 Color changes of MO and OG solutions before and 48 h after adsorption onto **3.1** and **3.2** at pH = 7.20

Figure 3.8 Photographs of sorbents on adsorption of MO and OG after 48h

Figure 3.9 Freundlich isotherms for the adsorption of MO onto (a) **3.1** and (b) **3.2** [pH 7.20, temperature: 25°C]

Figure 3.10 Freundlich isotherms for the adsorption of OG onto (a) **3.1** and (b) **3.2** [pH 7.20, temperature: 25°C]

Figure 3.11 Desorption of MO and OG from dye loaded networks in water at pH = 12.0 (A) MO loaded **3.1** (B) OG loaded **3.1**(C) MO loaded **3.2** and (D) OG loaded **3.2**

Figure 3.12 A proposed mechanism of adsorption-desorption of azo dye pollutants onto the hybrid sorbents

Figure 4.1. 3-aminopyridine rich novolac-based network **4.1**

Figure 4.2 FTIR spectrum of **4.1**

Figure 4.3 solid state ^{13}C NMR spectrum of **4.1**

Figure 4.4 FESEM images of **4.1**.

Figure 4.5 TGA Thermogram of **4.1**

Figure 4.6 TGA Thermogram of **4.1**

Figure 4.7. UV-vis spectral changes of the solutions of (a) MO ($1.82 \times 10^{-4}\text{M}$; pH 7.20), (b) O II ($4.28 \times 10^{-4}\text{M}$; pH 10.96) and (c) OG ($3.31 \times 10^{-4}\text{M}$; pH 2.30) as a function of adsorption time for **4.1**; (temperature: 25°C).

Figure 4.8 Time profile for azo-dye adsorption on **4.1** at different pH (a) MO adsorption ($C_0 = 60 \text{ mg L}^{-1}$; $1.82 \times 10^{-4} \text{ M}$); (b) O II adsorption ($C_0 = 150 \text{ mg L}^{-1}$; $4.28 \times 10^{-4} \text{ M}$); (c) OG adsorption ($C_0 = 150 \text{ mg L}^{-1}$; $3.31 \times 10^{-4} \text{ M}$).

Figure 4.9 Effect of solution pH on the adsorption capacity of **4.1** for 48h of contact time [MO adsorption ($C_0 = 60 \text{ mg L}^{-1}$; $1.82 \times 10^{-4} \text{ M}$); O II adsorption ($C_0 = 150 \text{ mg L}^{-1}$; $4.28 \times 10^{-4} \text{ M}$); OG adsorption ($C_0 = 150 \text{ mg L}^{-1}$; $3.31 \times 10^{-4} \text{ M}$); 25°C].

Figure 4.10 The fittings of adsorption isotherm data to Freundlich model for the adsorption of MO onto **4.1** at (a) pH 2.30, (b) pH 7.20 and (c) pH 10.96 at 25°C

Figure 4.11 The fittings of adsorption isotherm data to Freundlich model for the adsorption of OII onto **4.1** at (a) pH 2.30, (b) pH 7.20 and (c) pH 10.96 at 25°C

Figure 4.12 The fittings of adsorption isotherm data to Freundlich model for the adsorption of OG onto **4.1** at (a) pH 2.30, (b) pH 7.20 and (c) pH 10.96 at 25°C

Figure 4.13 Photo showing color change of **4.1** on adsorption of azo dyes for 48h.

Figure 4.14 Photo image of adsorptive separation of azo dyes using sorbent **4.1** (a) MO adsorption ($C_0 = 60 \text{ mg L}^{-1}$; pH 7.20) (b) O II adsorption ($C_0 = 150 \text{ mg L}^{-1}$; pH 10.96) (c) OG adsorption ($C_0 = 150 \text{ mg L}^{-1}$; pH 2.30); 25°C .

Figure 4.15 A proposed mechanism of adsorption: Attractive interactions (electrostatic attraction, hydrogen bonds and π - π stacking) occurring within the network polymer matrix.

Figure 5.1 The structures of functionalized novolac resin **5.1** and its hybrid with AgNPs **5.3**

Figure 5.2. FTIR spectrum of **5.1**

Figure 5.3. ^1H NMR spectrum of **5.1**

Figure 5.4. Digital optical images showing color of the solutions of (a) **5.2** and (b) **5.3** in DMSO-water (20:1, v/v) ($[\text{AgNO}_3] = 6.38 \text{ mM}$).

Figure 5.5. UV-vis spectra of (a) **5.2** and (b) **5.3** in DMSO-water (20:1, v/v) ($[\text{AgNO}_3] = 6.38 \text{ mM}$).

Figure 5.6. TEM image and corresponding histogram of **5.3**

Figure 5.7. Images showing inhibitory zones of **5.3** hybrid against (a) *Staphylococcus aureus* (b) *Bacillus subtilis* and (c) *Escherichia coli* on agar plates.

Figure 5.8. Growth curves of representative pathogenic microorganisms exposed to (a) **5.1** and (b) **5.3**. Amount of hybrid **5.3** used in this assay as the obtained MIC values.

Figure 5.9. Bar graph showing the effect of **5.3** and **5.1** on the release of cytoplasmic material. The bacteria were incubated with a fixed concentration ($4.0 \mu\text{L}$) at 30°C for 12 h. The change in the absorbance at 260 nm was monitored. Data are the mean of triplicates \pm S.E.