

List of figures

| Figure no. with the heading | Page no. |
|--|----------|
| Fig. 2.1 Options for reconstruction of degraded ecosystem (Bradshaw, 1996) | 16 |
| Fig. 2.2. As time passes, grass and legume covers enhance yield of organic matter and nitrogen for the fast-growing trees and desire multipurpose trees (MPT), which gradually mature and develop forest (Modified after Burger and Zipper 2002). | 32 |
| Figure 2.3. P transformations with time. P_{Ca} = calcium phosphates, P_T = total P, P_o = P bound to organic matter | 56 |
| Fig 3.1 Schematic diagram of the experimental protocol | 60 |
| Fig 3.2 Location of Jharia Coalfields | 62 |
| Fig. 3.9 SEDEX scheme for different P fraction (Ruttenberg, 1992) | 84 |
| Fig. 4.1. Temperature and Humidity in Jharia from April 2012 to December 2013. | 93 |
| Fig.4.2. Total rainfall (mm) in Jharia from April 20014 to December 2015. | 93 |
| Figure 4.3 Randomised Block Design of the Plots. A, B, C, D and E are the five individual plant species namely <i>Dendrocalamus strictus</i> , <i>Phullanthus emblica</i> , <i>Saraca asoca</i> , <i>Ficus religiosa</i> and <i>Azadirachta indica</i> . | 95 |
| Fig. 4.3 Percent survival of the total number of plants in various plots | 107 |
| Fig. 4.4 Percent survival of the total number of individual plant species in all the plots taken together | 107 |
| Fig. 4.5 Initial (2 months old individuals) and final (two years old) root and shoot biomass (dry weight, g) in control and treatment plots (period of study 2009-2011) | 108 |
| Fig. 4.6 Initial (2 months old individuals) and final (two years old) height (cm) in control and treatment plots from 2009 to 2011 | 108 |
| Fig. 4.7. Phosphorus fractions found in the overburden dump samples | 116 |
| Fig. 4.8. XRD report of the overburden dump samples | 116 |
| Fig. 4.9 Mass loss of litter of the five species with time in the four plots | 120 |