

List of tables

Table no. with the heading	Page no.
Table 1.1 State wise Estimated Reserves of Coal in India as on 31.03.2010 (Office of Coal Controller, Ministry of Coal)	3
Table 2.1 Ecosystem characteristics for consideration as ecological restoration objectives (Cooke and Jhonson, 2002)	20
Table 2.2 Selection criteria of plant species based on spoil characteristics and future land use pattern (Maiti, 2006).	30
Table 2.3 Suggested multipurpose tree (MPTs) for revegetation of mine spoil (CPCB 2000)	34
Table 2.4 Spore density in different rhizosphere of host plant in overburden dumps (Maiti and Shee, 2003).	48
Table 2.5 Litter accumulation and decomposition rate in a reclaimed coal mine overburden dumps (all weights in g dry wt/m ²) (Mukhopadhyay and Maiti, 2010).	53
Table 3.3. Materials: list of materials used is enlisted	66
Table 3.4. Detail of the control and treatment plots	74
Table 3.5. Methods used for various physicochemical estimation	76
Table 4.1 Physico-chemical parameters of the coal mine spoil, agricultural soil, and cow dung manure	99
Table 4.2. Physico-chemical parameters of the control and treatment plots in three years of study (DTPA extractable metals are depicted)	101
Table 4.3. Physico-chemical parameters of the control and treatment plots in three years of study 2013, 2014, and 2015	102

Table 4.4 Determination of soil pollution index in the overburden dump of Bastacola (Alloway, 1990)	103
Table 4.7. Percentage increase in root biomass, shoot biomass, circumference, and height of the plants from Aug'13 to Aug'2015 under the control and treatment plots	110
Table 4.8 H: D ratio, Relative Growth Rate in height (RGRH, cm/month) and circumference (RGRC, cm/month), annual increment in height (cm) and circumference (cm), between Aug'13 and Aug '15 in control and experimental plots	114
Table 4.9 Initial chemical composition of leaf litter of the five plant species	119
Table 4.10 Daily decay rate coefficient of litter decomposition (k_d), time required in days for 50% (t_{50}) and 99% (t_{99}) weight loss and N (k_N), P (k_P), and K (k_K) losses of 5 plant species in 4 plots	121
Table 4.11. F values for k_d , k_N , k_P and k_K at 5 % level of significance	123
Table 4.12 Seasonal variation in DHA activity ($\mu\text{g/g/hr}$) in the rhizosphere of five plant species along with their F values and outcomes of DMRT	131
Table 4.13 Seasonal variation in catalase activity ($\text{ml KMnO}_4/\text{g}$) in the litter of five plant species along with their F values	138
Table 4.14 Percent root infection of VAM with the five plant species	139
Table 5.1 Ethnomedicinal uses of different parts of <i>F. religiosa</i>	144