

List of Figures

- Fig. 1.1** Formation of 1D, 2D and 3D network.
- Fig. 1.2** Some representative linkers for coordination polymers.
- Fig. 1.3** Arrangement of interactions contributing in coordination polymers.
- Fig. 1.4** Zn(II) 1D chain.
- Fig. 1.5** (a) Fragment of a layer displaying the $[\text{Fe}(\mu\text{-tvp})]_n^{2n+}$ chains connected through $\{[\text{Hg}^{\text{II}}(\text{SCN})_3]_2(\mu\text{-L})\}^{2-}$ units. (b) View down the [001] direction of the connection between two chains (note that this connection contains another relevant inversion center). (c) Stack of two consecutive layers. Coordination centers: $[\text{FeN}_6]$ (orange), $[\text{HgS}_3\text{N}]$ (pink).
- Fig. 1.6** Structure of the CP showing (a) the coordination environments of the Fe centers, C black, O red, F green, Fe1 blue, Fe2 pink; Fe2 is located on a crystallographic threefold axis. (b) One of two (10,3)-a networks. (c) Schematic representation of the two independent (10,3)-a nets; spheres represent Fe centers (nodes), and connections represent bridging fan anions. (d) Space-filling representation of the two anionic networks (green and purple).
- Fig. 2.1** The trinuclear synthon in complex **1** that gives origin to 2D corrugated layer.
- Fig. 2.2** Coordination sphere of Pb1 and Pb2 in **1**. Of the latter the crystallographic disorder is shown.
- Fig. 2.3** The 2D network in the *ab*-plane (for clarity phen ligands not shown) for **1**. The light blue spheres indicated the disordered Pb(2) ions.
- Fig. 2.4** Perspective views of the interdigitated 2D layers with *o*-phen molecules interacting through $\pi - \pi$ interactions in **1**.

- Fig. 2.5** ORTEP drawing (ellipsoids at 40% probability) of the coordination sphere of the lead(II) ion in complex **2**. The Pb-O(1)^{iv} and Pb-O(1)^v are long distances of 2.931(5) Å.
- Fig. 2.6** The one-dimensional Pb chains connected by the bridging carboxylates groups in **2**.
- Fig. 2.7** The inorganic-organic hybrid structure 3D structure of **2**.
- Fig. 2.8** Emission spectra of (a) O-phen, (b) complex **1**.
- Fig. 2.9** Emission spectra of (a) ndc, (b) complex **2**.
- Fig. 2.10** IR spectrum of complex **1**.
- Fig. 2.11** IR spectrum of complex **2**.
- Fig. 3.1** Coordination sphere of Zn ion in compound **3**.
- Fig. 3.2** Zig-zag coordination polymer of **3**.
- Fig. 3.3** The coordination polymers connected by π - π interactions occurring between bipy rings.
- Fig. 3.4** Coordination sphere of the two independent cadmium atoms in **4**.
- Fig. 3.5** The couple of coordination polymers formed by Cd1 and Cd2 ions in compound **4**. The H-bond connections of O1w with carboxylate oxygens of the chains are also shown.
- Fig. 3.6** Coordination sphere of cadmium atom in **5**. The bridging nitrate is located on a crystallographic two-fold axis.
- Fig. 3.7** Coordination polymer of compound **5**.
- Fig. 3.8** Crystal packing viewed down axis *c* of compound **5** showing two coordination polymers and the channels filled by lattice water molecules and uncoordinated nitrate anions.
- Fig. 3.9** Emission spectrum of 5-nitro-isophthalic acid, H₂NIP ($\lambda_{\text{Ex}} = 280$ nm).
- Fig. 3.10** Emission spectrum of **3** ($\lambda_{\text{Ex}} = 295$ nm).

- Fig. 3.11** Emission spectrum of **4** ($\lambda_{\text{Ex}} = 280$ nm).
- Fig. 3.12** Emission spectrum of 1,4-benzenediacrylic acid, H₂BDAC ($\lambda_{\text{Ex}} = 280$ nm).
- Fig. 3.13** Emission spectrum of **5** ($\lambda_{\text{Ex}} = 243$ nm).
- Fig. 3.14** IR spectrum of complex **3**, **4** and **5**.
- Fig. 4.1** The centrosymmetric trinuclear unit in compound **6**.
- Fig. 4.2** The 2D architecture of compound **6**.
- Fig. 4.3** 2D layers viewed down axis *a* with indications of π - π interaction (in green). Lattice water molecules indicated as big red sphere also connect the polymeric layers through H-bonds.
- Fig. 4.4** A symmetric unit of compound **7** with label scheme of N and O atoms.
- Fig. 4.5** Crystal packing of compound **7** with indication of H-bonds among the Mn(H₂O)₄(bipy)]_n polymers, the dicarboxylate anions and the lattice water molecule.
- Fig. 4.6** The centrosymmetric dinuclear entity in compound **8** with indication of intramolecular H-bonds.
- Fig. 4.7** H-bonds connecting dinuclear complexes to form a polymeric chain along axis *a*.
- Fig. 4.8** Emission spectrum of 1,4-phenylenediacetic, H₂phdac acid ($\lambda_{\text{Ex}} = 275$ nm).
- Fig. 4.9** Emission spectrum of 1,10-phenethroline, phen ($\lambda_{\text{Ex}} = 243$ nm).
- Fig. 4.10** Emission spectrum of **6** ($\lambda_{\text{Ex}} = 280$ nm).
- Fig. 4.11** Emission spectrum of 4-nitrophthalic acid, H₂nph ($\lambda_{\text{Ex}} = 270$ nm).
- Fig. 4.12** Emission spectrum of **8** ($\lambda_{\text{Ex}} = 300$ nm).
- Fig. 4.13** FT-IR Spectrum of **6**.
- Fig. 4.14** FT-IR Spectrum of **7**.
- Fig. 5.1** Coordination environment and molecular view of **9** with atom numbering scheme. Solvent water molecules have been omitted for clarity.
- Fig. 5.2** Propagation of the 1D polymeric chain along (0 0 1) direction.

- Fig. 5.3** (a) Formation of the 2D layer architecture through O–H···O hydrogen bonding interactions in **9**. (b) Generation of the 3D architecture (Parallel view) in **9** through weak C–H···O hydrogen bonding interactions.
- Fig. 5.4** The central projection of the 3D architecture in **9**.
- Fig. 5.5** (a) Formation of supramolecular layered assembly generated through C–H··· π interactions in **9**. (b) Perspective view metal··· π interaction in building layer network in (0 1 1) plane in **9**.
- Fig. 5.6** Coordination environment and molecular view of **10** with atom numbering scheme.
- Fig. 5.7** (a) Formation of polymeric network in (0 1 1) plane in **10**. (b) Generation of 2D network in (0 1 1) plane through O–H···O hydrogen bonding interactions in **10**. (c) Generation of 2D network in (0 1 1) plane through C–H···O hydrogen bonding interactions in **10**.
- Fig. 5.8** (a) Formation of supramolecular layered assembly in (1 0 1) plane through C–H··· π interactions in **10**. (b) Formation of 2D layer network in (0 1 1) plane in **10** generated through metal··· π interaction.
- Fig. 5.9** Emission spectrum of 4, 4'-trimethylene dipyramine, bpp ($\lambda_{\text{EX}} = 280$ nm).
- Fig. 5.10** Emission spectrum of **9** ($\lambda_{\text{EX}} = 350$ nm).
- Fig. 5.11** Emission spectrum of 5-Sulfosalicylic acid, H₃SSA ($\lambda_{\text{EX}} = 280$ nm).
- Fig. 5.12** Emission spectrum of **10** ($\lambda_{\text{EX}} = 280$ nm).
- Fig. 5.13** IR spectrum of complex **9**.
- Fig. 5.14** IR spectrum of complex **10**.
- Fig. 6.1** ORTEP drawing (50% probability level) of the crystallographic independent unit in complex **11**. Coordination bond distances (Å): Pb–N(1) 2.556(4), Pb–O(3')2.689(4), Pb–O(1) 2.596(4), Pb–O(4') 2.334(4), Pb–O(2) 2.342(3).
- Fig. 6.2** The 1D coordination polymer in compound **11**.

- Fig. 6.3** ORTEP drawing (50% probability level) of the crystallographic independent unit in complex **12** with indication of the coordination sphere around Pb atom (primed atoms at 1-x, 2-y, 1-z).
- Fig. 6.4** A single 1D polymeric chain in **12** built by centrosymmetrical $[\text{Pb}_2(\text{dbsf})_2]$ units connected by bipy spacers.
- Fig. 6.5** The interpenetrated 1D coordination polymer in compound **12** as ball and stick and space-filling representation.
- Fig. 6.6** Schematic representation of possible self-interpenetrating 1D-1D polymers.
- Fig. 6.7** TG analysis of **11** in the temperature range 25-600°C with a heating rate of 10°C/min under N_2 atmosphere.
- Fig. 6.8** TG analysis of **12** in the temperature range 0-800°C with a heating rate of 10°C/min under N_2 atmosphere.