

Bibliography

- Abbas, T., Nadeem, M. A., Tanveer, A., & Chauhan, B. S. (2017). Can hormesis of plant-released phytotoxins be used to boost and sustain crop production? *Crop Protection*, 93, 69–76. doi:10.1016/j.cropro.2016.11.020
- Achatz, M., Morris, E. K., Müller, F., Hilker, M., & Rillig, M. C. (2014). Soil hypha-mediated movement of allelochemicals: arbuscular mycorrhizae extend the bioactive zone of juglone. *Functional Ecology*, 28(4), 1020–1029. doi:10.1111/1365-2435.12208
- Adler, M. J., & Chase, C. A. (2007). Comparison of the Allelopathic Potential of Leguminous Summer Cover Crops: Cowpea, Sunn Hemp, and Velvetbean. *Hort Science*.
- Akhter, J., Murray, R., Mahmood, K., Malik, K. A., & Ahmed, S. (2004). Improvement of degraded physical properties of a saline-sodic soil by reclamation with kallar grass (*Leptochloa fusca*). *Plant and Soil*, 258(1), 207–216. doi:10.1023/B:PLSO.0000016551.08880.6b
- Ames, B. N. (1979). Identifying environmental chemicals causing mutations and cancer. *Science*, 204(4393), 587–593. doi:10.1126/science.373122
- Andrioli NB, Soloneski S, Laramendy ML, Mudry MD (2012) Cytogenetic and microtubule array effects of the zineb- containing commercial fungicide formulation Azzurro® on meristematic root cells of *Allium cepa* L. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 742(1-2): 48-53.
- Aronson, J. A., & Whitehead, E. E. (1989). HALOPH: a data base of salt tolerant plants of the world.
- Babikova, Z., Gilbert, L., Bruce, T. J. A., Birkett, M., Caulfield, J. C., Woodcock, C., ... Johnson, D. (2013). Underground signals carried through common mycelial networks warn neighbouring plants of aphid attack. *Ecology Letters*, 16(7), 835–843. doi:10.1111/ele.12115
- Baerson, S. R., Sánchez-Moreiras, A., Pedrol-Bonjoch, N., Schulz, M., Kagan, I. A., Agarwal, A. K., ... Duke, S. O. (2005). Detoxification and transcriptome response in *Arabidopsis* seedlings exposed to the allelochemical benzoxazolin-2(3H)-one. *The Journal of Biological Chemistry*, 280(23), 21867–21881. doi:10.1074/jbc.M500694200
- Bais, H. P., Park, S.-W., Weir, T. L., Callaway, R. M., & Vivanco, J. M. (2004). How plants communicate using the underground information superhighway. *Trends in Plant Science*, 9(1), 26–32. doi:10.1016/j.tplants.2003.11.008

- Banerjee, P., & Talapatra, S. N. (2015). Assessment of medicinal tree diversity in the Chintamoni Kar Bird Sanctuary (CKBS), Kolkata, India and prediction of antimutagenic phytochemicals by using software. *Int. J. Adv. Res.*, 3(7), 225–243.
- Barto, E. K., Weidenhamer, J. D., Cipollini, D., & Rillig, M. C. (2012). Fungal superhighways: do common mycorrhizal networks enhance below ground communication? *Trends in Plant Science*, 17(11), 633–637. doi:10.1016/j.tplants.2012.06.007
- Batish, D R, Singh, H. P., Pandher, J. K., & Kohli, R. K. (2005). Invasive potential of the weed *Parthenium hysterophorus*-the role of allelopathy. *Plant Protection and Plant Health in Europe: Introduction and Spread of Invasive Species*, Held at Humboldt University, Berlin, Germany, 9–11.
- Batish, Daizy R., Lavanya, K., Singh, H. P., & Kohli, R. K. (2007). Phenolic allelochemicals released by *Chenopodium murale* affect the growth, nodulation and macromolecule content in chickpea and pea. *Plant Growth Regulation*, 51(2), 119–128. doi:10.1007/s10725-006-9153-z
- Baziramakenga, R., Leroux, G. D., & Simard, R. R. (1995). Effects of benzoic and cinnamic acids on membrane permeability of soybean roots. *Journal of Chemical Ecology*, 21(9), 1271–1285. doi:10.1007/BF02027561
- Benyas, E., Hassanpouraghdam, M. B., Zehtabsalmasi, S., & Oskooei, O. K. (2010). Allelopathic effects of *Xanthium strumarium L.* shoot aqueous extract on germination, seedling growth and chlorophyll content of lentil (*Lens culinaris Medic.*). *Romanian Biotechnological Letters*, 15(3), 5223–5228.
- Bever, J. D., Westover, K. M., & Antonovics, J. (1997). Incorporating the Soil Community into Plant Population Dynamics: The Utility of the Feedback Approach. *The Journal of Ecology*, 85(5), 561. doi:10.2307/2960528
- Bhadoria, P. (2010). Allelopathy: A Natural Way towards Weed Management. *American Journal of Experimental Agriculture*, 1(1), 7–20. doi:10.9734/AJEA/2011/002
- Biswas, A. K., & Choudhuri, M. A. (1978). Differential Behaviour of the Flag Leaf of Intact Rice Plant during Ageing. *Biochemie Und Physiologie Der Pflanzen*, 173(3), 220–228. doi:10.1016/S0015-3796(17)30485-7
- Bjeldanes, L. F., & Chang, G. W. (1977). Mutagenic activity of quercetin and related compounds. *Science*, 197(4303), 577–578. doi:10.1126/science.327550
- Blackburn, T. M., Cassey, P., Duncan, R. P., Evans, K. L., & Gaston, K. J. (2004). Avian extinction and mammalian introductions on oceanic islands. *Science*, 305(5692), 1955–1958. doi:10.1126/science.1101617

Blum, U. (1998). :{unav}. Springer Science and Business Media LLC. doi:10.1023/a:1022394203540

Borges de Melo, E., Ataide Martins, J. P., Marinho Jorge, T. C., Friozi, M. C., & Castro Ferreira, M. M. (2010). Multivariate QSAR study on the antimutagenic activity of flavonoids against 3-NFA on *Salmonella typhimurium* TA98. *European Journal of Medicinal Chemistry*, 45(10), 4562–4569. doi:10.1016/j.ejmech.2010.07.017

Börner, H. (1960). Liberation of organic substances from higher plants and their role in the soil sickness problem. *The Botanical Review*, 26(3), 393–424. doi:10.1007/BF02860808

Brown, J. P., & Brown, R. J. (1976). Mutagenesis by 9,10-anthraquinone derivatives and related compounds in *Salmonella typhimurium*. *Mutation Research/Genetic Toxicology*, 40(3), 203–224. doi:10.1016/0165-1218(76)90046-X

Callaway, R. M. (2002). The detection of neighbors by plants. *Trends in Ecology & Evolution*, 17(3), 104–105. doi:10.1016/S0169-5347(01)02438-7

Callaway, R. M., Cipollini, D., Barto, K., Thelen, G. C., Hallett, S. G., Prati, D., Klironomos, J. (2008). Novel weapons: invasive plant suppresses fungal mutualists in America but not in its native Europe. *Ecology*, 89(4), 1043–1055. doi:10.1890/07-0370.1

Chen, B.-M., Peng, S.-L., & Ni, G.-Y. (2009). Effects of the invasive plant *Mikania micrantha* H.B.K. on soil nitrogen availability through allelopathy in South China. *Biological Invasions*, 11(6), 1291–1299. doi:10.1007/s10530-008-9336-9

Chen, Y., Paetz, C., & Schneider, B. (2018). Precursor-Directed Biosynthesis of Phenylbenzoisoquinolindione Alkaloids and the Discovery of a Phenylphenalenone-Based Plant Defense Mechanism. *Journal of Natural Products*, 81(4), 879–884. doi:10.1021/acs.jnatprod.7b00885

Choudhuri, M. A., & Chatterjee, S. K. (1970). Seasonal changes in the levels of some cellular components in the abscission zone of *Coleus* leaves of different ages. *Annals of Botany*, 34(2), 275–287.

Coolbear, P., Francis, A., & Grierson, D. (1984). The effect of low temperature pre-sowing treatment on the germination performance and membrane integrity of artificially aged tomato seeds. *Journal of Experimental Botany*, 35(11), 1609–1617.

Croteau, R. B., Bohlmann, J., Crock, J. E., & Steele, C. L. (2002). Sesquiterpene synthases from grand fir (*Abies grandis*), and methods of use. Washington, DC.

- Cummings, J. A., Parker, I. M., & Gilbert, G. S. (2012). Allelopathy: a tool for weed management in forest restoration. *Plant Ecology*, 213(12), 1975–1989. doi:10.1007/s11258-012-0154-x
- Dabaghao, P. M., & Shankarnarayan, K. A. (1973). *The grass cover of India. The Grass Cover of India.*
- Dallwitz, M. J. (1993). *Applications and documentation of the DELTA System.*
- Das, V. R., & Vats, S. K. (1993). A Himalayan monsoon location exhibiting unusually high preponderance of C4 grasses. *Photosynthetica*, 28(1), 91–97.
- Datta, S. C., & Biswas, K. K. (1979). Autecological studies on weeds of West Bengal. VIII *Alternanthera sessilis* (L.) DC. *Bulletin of the Botanical Society of Bengal*, 33(1), 5–26.
- De, G. C. (1983). *Parthenium hysterophorus L. A national hazard. Pesticides*, 17, 31–36.
- De Galicia, X. (2007). *Plantas invasoras de Galicia: biología, distribución e métodos de control.*
- Dolling, A., Zackrisson, O., & Nilsson, M. C. (1994). Seasonal variation in phytotoxicity of bracken (*Pteridium aquilinum* L. Kuhn). *Journal of Chemical Ecology*, 20(12), 3163–3172. doi:10.1007/BF02033718
- Dyer, L. A., Dodson, C. D., Stireman, J. O., Tobler, M. A., Smilanich, A. M., Fincher, R. M., & Letourneau, D. K. (2003). Synergistic effects of three *Piper* amides on generalist and specialist herbivores. *Journal of Chemical Ecology*, 29(11), 2499–2514. doi:10.1023/a:1026310001958
- Edenharder, R., & Tang, X. (1997). Inhibition of the mutagenicity of 2-nitrofluorene, 3-nitrofluoranthene and 1-nitropyrene by flavonoids, coumarins, quinones and other phenolic compounds. *Food and Chemical Toxicology*, 35(3-4), 357–372. doi:10.1016/s0278-6915(97)00125-7
- Einhellig, F. A. (1996). Interactions involving allelopathy in cropping systems. *Agronomy Journal*, 88(6), 886. doi:10.2134/agronj1996.00021962003600060007x
- El-Darier, S. M., Abdelaziz, H. A., & Zein El-Dien, M. H. (2014). Effect of soil type on the allelopathic activity of *Medicago sativa* L. residues in *Vicia faba* L. agroecosystems. *Journal of Taibah University for Science*, 8(2), 84–89. doi:10.1016/j.jtusci.2014.01.001
- Ellstrand, N. C., & Schierenbeck, K. A. (2000). Hybridization as a stimulus for the evolution of invasiveness in plants? *Proceedings of the National Academy of Sciences of the United States of America*, 97(13), 7043–7050. doi:10.1073/pnas.97.13.7043

- Elton, C. S. (1958). The Ecology of Invasions by Animals and Plants. *Methuen and Co Ltd. London. England.*
- Evenari, M. (1949). Germination inhibitors. *The Botanical Review*, 15(3), 153–194. doi:10.1007/BF02861721
- Fargione, J. E., & Tilman, D. (2005). Diversity decreases invasion via both sampling and complementarity effects. *Ecology Letters*, 8(6), 604–611. doi:10.1111/j.1461-0248.2005.00753.x
- Fick, G. N., & Qualset, C. O. (1975). Genetic control of endosperm amylase activity and gibberellic Acid responses in standard-height and short-statured wheats. *Proceedings of the National Academy of Sciences of the United States of America*, 72(3), 892–895. doi:10.1073/pnas.72.3.892
- Firn, R. D., & Jones, C. G. (2000). The evolution of secondary metabolism - a unifying model. *Molecular Microbiology*, 37(5), 989–994. doi:10.1046/j.1365-2958.2000.02098.x
- Fitter, A. (2003). Ecology. Making allelopathy respectable. *Science*, 301(5638), 1337–1338. doi:10.1126/science.1089291
- Fridley, J. D., Stachowicz, J. J., Naeem, S., Sax, D. F., Seabloom, E. W., Smith, M. D., Von Holle, B. (2007). The invasion paradox: reconciling pattern and process in species invasions. *Ecology*, 88(1), 3–17. doi:10.1890/0012-9658(2007)88[3:tiprpa]2.0.co;2
- Fu, Y. H., Quan, W. X., Li, C. C., Qian, C. Y., Tang, F. H., & Chen, X. J. (2019). Allelopathic effects of phenolic acids on seedling growth and photosynthesis in *Rhododendron delavayi* Franch. *Photosynthetica*, 57(2), 377–387. doi:10.32615/ps.2019.045
- Gaertner, M., Den Breeyen, A., Cang Hui, & Richardson, D. M. (2009). Impacts of alien plant invasions on species richness in Mediterranean-type ecosystems: a meta-analysis. *Progress in Physical Geography: Earth and Environment*, 33(3), 319–338. doi:10.1177/030913309341607
- Ghayal, N. A., Dhumal, K. N., Deshpande, N. R., Ruikar, A. D., & Phalgune, U. D. (2011). Phytotoxic effects of leaf leachates of an invasive weed *Cassia uniflora* and characterization of its allelochemical. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 2(4), 524–534.
- Gnanavel, I., & Natarajan, S. K. (2013). *Parthenium hysterophorus L.*: a major threat to natural and agro eco-systems in India. *International Journal of Agriculture, Environment and Biotechnology*, 6(2), 261–9.

- Golla, U., Gajam, P. K., & Raj B, S. S. (2013). The Effect of *Desmostachya bipinnata* (Linn.) Extract on Physiologically Altered Glycemic Status in Non-diabetic Rats. *Journal of Medical Sciences(Faisalabad)*, 13(3), 221–225. doi:10.3923/jms.2013.221.225
- Grümmer, G. (1955). Die gegenseitige Beeinflussung höherer Pflanzen: Allelopathie. *Die Gegenseitige Beeinflussung Höherer Pflanzen: Allelopathie*.
- Gupta, S., & Palit, D. (2014). Biosurveillance of wetlands in Eastern India (Birbhum, West Bengal) for wise use. *International Journal of Science, Environment and Technology*, 3(6), 2136–2144.
- Gupta, S. R., Jangra, R., & Dagar, J. C. (2015). Carbon pools and fluxes in grassland systems on sodic soils of northern india. In A. K. Singh, J. C. Dagar, A. Arunachalam, G. R, & K. N. Shelat (Eds.), *Climate Change Modelling, Planning And Policy For Agriculture* (pp. 119–140). New Delhi: Springer India. doi:10.1007/978-81-322-2157-9_14
- Hakoo, M. L. (1963). A diploid *Parthenium* in Jammu. *Current Science*, 32(6), 273–273.
- Halder, S., & Gupta, K. (1982). On the mechanism of sunflower seed deterioration under low and high RH. *Seed Science and Technology*, 10(2), 267–276.
- Hansen, K., Mika, S., Schroeter, T., Sutter, A., ter Laak, A., Steger-Hartmann, T., ... Müller, K.-R. (2009). Benchmark data set for in silico prediction of Ames mutagenicity. *Journal of Chemical Information and Modeling*, 49(9), 2077–2081. doi:10.1021/ci900161g
- Haque M, Bhakat RK , Bhattacharjee A (2018) Evaluation of relative allelopathic potential of three invasive weeds of West Bengal using their leaf extracts on cells of *Allium cepa* root tips. *Indian Biologist* 50(2): 13-19.
- Hegde, B. A., & Patil, T. M. (1982). Effect of salt stress on the structure and carbon flow mechanism in a noxious weed *Parthenium hysterophorus* L. *Weed Research*, 22(1), 51–56. doi:10.1111/j.1365-3180.1982.tb00143.x
- Hegde, M. M., Lakshman, K., Girija, K., Kumar, B., & Lakshmiprasanna, V. (2010). Assessment of antidiarrhoeal activity of *Desmostachya bipinnata* L.(Poaceae) root extracts. *Boletín Latinoamericano y Del Caribe de Plantas Medicinales y Aromáticas*, 9(4), 312–318.
- Hejda, M., Pyšek, P., & Jarošík, V. (2009). Impact of invasive plants on the species richness, diversity and composition of invaded communities. *Journal of Ecology*, 97(3), 393–403. doi:10.1111/j.1365-2745.2009.01480.x
- Herrero O, Martín JP, Freire PF, López LC, Peropadre A, Hazen MJ (2012) Toxicological evaluation of three contaminants of emerging concern by use of the *Allium cepa* test.

Mutation Research/Genetic Toxicology and Environmental Mutagenesis, 743(1-2): 20-24.

Hifnawy, M. S., El-Hyatmy, Y. Y., Kenawy, S. A., Yossef, A. K., & Awaad, A. S. (1999).

Carbohydrate, lipid, protein and amino acid contents of *Chenopodium murale*, *Cyperus alopecuroides*, *Desmostachya bipinnata* and *Tamaric nilotica* as allergic plants. *Bull Fac Cairo Univ*, 37, 99–106.

Holopainen, J. K., Virjamo, V., Ghimire, R. P., Blande, J. D., Julkunen-Tiitto, R., & Kivimäenpää, M. (2018). Climate change effects on secondary compounds of forest trees in the northern hemisphere. *Frontiers in Plant Science*, 9, 1445. doi:10.3389/fpls.2018.01445

HR Bode. (1940). *Planta. Planta*.

Hulme, P. E., Pysek, P., Nentwig, W., & Vilà, M. (2009). Ecology. Will threat of biological invasions unite the European Union? *Science*, 324(5923), 40–41. doi:10.1126/science.1171111

Hunt, S., Smith, A. M., & Woolhouse, H. W. (1987). Evidence for a light-dependent system for reassimilation of photorespiratory CO₂, which does not include a C 4 cycle, in the C 3-C 4 intermediate species *Moricandia arvensis*. *Planta*, 171(2), 227–234. doi:10.1007/BF00391098

Hussain, F., & Rashid, A. (1989). A checklist to the monocotyledonous weeds of Pakistan I. Family Poaceae. *Sarhad Journal of Agriculture*, 5(2), 183–197.

Hussain, M. I., & Reigosa, M. J. (2011). Allelochemical stress inhibits growth, leaf water relations, PSII photochemistry, non-photochemical fluorescence quenching, and heat energy dissipation in three C₃ perennial species. *Journal of Experimental Botany*, 62(13), 4533–4545. doi:10.1093/jxb/err161

Inderjit, C. H., & Nishimura, H. (1999). Plant phenolics and terpenoids: transformation, degradation, and potential for allelopathic interactions. *Principles and Practices in Plant Ecology: Allelochemical Interactions* (Inderjit, KMM et Al., Eds), 255–266.

Inderjit, & Dakshini, K. M. M. (1998). Allelopathic interference of chickweed, *Stellaria media* with seedling growth of wheat (*Triticum aestivum*). *Canadian Journal of Botany*, 76(7), 1317–1321. doi:10.1139/b98-159

Inderjit, & Weiner, J. (2001). Plant allelochemical interference or soil chemical ecology? *Perspectives in Plant Ecology, Evolution and Systematics*, 4(1), 3–12. doi:10.1078/1433-8319-00011

- Iannacone JO, Onofre RC, Huanqui OS, Giraldo JA, Mamani NP, Miglio MC, Alvariño LF (2007) Environmental risk assessment of the insecticide metamidophos in bioassays with four non-target aquatic organisms. *Agricultura Tecnica*, 67(2): 126-138.
- Indian Council of Agricultural Research, New Delhi, India. Reinhardt P, Causa M, Marian CM, Hess BA (1996) *Physical Review*, B54: 14812-14821.
- International Seed Testing Association. ISTA. (1996). International rules for seed testing. *Seed Sci. Technol.*, 27.
- Ishii-Iwamoto, E. L., Abraham, D., Sert, M. A., Bonato, C. M., Kelmer-Bracht, A. M., & Bracht, A. (2006). Mitochondria as a site of allelochemical action. *Allelopathy*, 267–284.
- Islam, A. K. M. M., & Kato-Noguchi, H. (2013). Allelopathic prospective of *Ricinus communis* and *Jatropha curcas* for bio-control of weeds. *Acta Agriculturae Scandinavica, Section B - Soil & Plant Science*, 63(8), 731–739. doi:10.1080/09064710.2013.865073
- Jacinthe, P.-A., Bills, J. S., & Tedesco, L. P. (2010). Size, activity and catabolic diversity of the soil microbial biomass in a wetland complex invaded by reed canary grass. *Plant and Soil*, 329(1-2), 227–238. doi:10.1007/s11104-009-0147-2
- Javaid, A., Bajwa, R., & Anjum, T. (2005). Biological Control of *Parthenium* II: Allelopathic effect of *Desmostachya bipinnata* on distribution and early seedling growth of *Parthenium hysterophorus* L. *International Journal of Biology and Biotechnology*.
- Jelenic, B. (1987). Allelopathy and allelochemicals from agrostemma githago as theoretical background of agrostemin production and use as a natural plant-growth bioregulator. *Agrostemin International Scientific Centre of Fertilizers*, 13–26.
- Jiang, X. L., Zhang, W. G., & Wang, G. (2007). Biodiversity effects on biomass production and invasion resistance in annual versus perennial plant communities. In D. L. Hawksworth & A. T. Bull (Eds.), *Plant Conservation and Biodiversity* (pp. 409–420). Dordrecht: Springer Netherlands. doi:10.1007/978-1-4020-6444-9_26
- Joshi, A., Bachheti, R. K., Sharma, A., & Mamgain, R. (2016). *Parthenium hysterophorus*. L. (Asteraceae) : A boon or curse? (A review). *Oriental Journal of Chemistry*, 32(3), 1283–1294. doi:10.13005/ojc/320302
- Kaminsky, R. A. Y., & Muller, W. H. (1977). The extraction of soil phytotoxins using a neutral edta solution. *Soil Science*, 124(4), 205–210. doi:10.1097/00010694-197710000-00003
- Kanchan, S., & Jayachandra. (1981). Effect of *Parthenium hysterophorus* on nitrogen-fixing and nitrifying bacteria. *Canadian Journal of Botany*, 59(2), 199–202. doi:10.1139/b81-030

- Kar, M., & Mishra, D. (1976). Catalase, peroxidase, and polyphenoloxidase activities during rice leaf senescence. *Plant physiology*, 57(2), 315-319.
- Kardol, P., Cornips, N. J., van Kempen, M. M. L., Bakx-Schotman, J. M. T., & van der Putten, W. H. (2007). Microbe-mediated plant-soil feedback causes historical contingency effects in plant community assembly. *Ecological Monographs*, 77(2), 147–162. doi:10.1890/06-0502
- Kaur, B., Gupta, S. R., & Singh, G. (2002). Bioamelioration of a sodic soil by silvopastoral systems in northwestern India. *Agroforestry Systems*, 54(1), 13–20.
- Kaur, M., Aggarwal, N. K., Kumar, V., & Dhiman, R. (2014). Effects and Management of *Parthenium hysterophorus*: A Weed of Global Significance. *International Scholarly Research Notices*, 2014, 368647. doi:10.1155/2014/368647
- Khan, A. A., & Faust, M. A. (1967). Effect of growth retardants on α -amylase production in germinating barley seed. *Physiologia Plantarum*, 20(3), 673-681.
- Khanh, T. D., Elzaawely, A. A., Chung, I. M., Ahn, J. K., & Xuan, T. D. (2007). Role of allelochemicals for weed management in rice. *Allelopathy Journal*.
- Kirtikar, K. R., & Basu, B. D. (1918). *Indian medicinal plants*. 4th vol. *International Book Distributors Book Sellers and Publishers, Dehradun*, 2688.
- Klironomos, J. N. (2002). Feedback with soil biota contributes to plant rarity and invasiveness in communities. *Nature*, 417(6884), 67–70. doi:10.1038/417067a
- Kohli, R. K., & Daizy, R. (1994). *Parthenium hysterophorus*-a review. *Research Bulletin of the Panjab University, Science*, 44(1), 105–149.
- Kourtev, P. S., Ehrenfeld, J. G., & Häggblom, M. (2003). Experimental analysis of the effect of exotic and native plant species on the structure and function of soil microbial communities. *Soil Biology and Biochemistry*, 35(7), 895–905. doi:10.1016/S0038-0717(03)00120-2
- Kumar, S., & Rohatgi, N. (1999). The role of invasive weeds in changing floristic diversity. *Annals of Forestry*, 7(1), 147–150.
- Lara-Nuñez, Romero-Romero, Ventura, J. L., Blancas, V., Anaya, A. L., & Cruz-Ortega, (2006). Allelochemical stress causes inhibition of growth and oxidative damage in *Lycopersicon esculentum* Mill. *Plant, Cell & Environment*, 29(11), 2009-2016.
- Leather, G. R. (1983). Weed control using allelopathic crop plants. *Journal of Chemical Ecology*, 9(8), 983–989. doi:10.1007/BF00982205

- Leme DM, Marin-Morales MA (2009) Allium cepa test in environmental monitoring: a review on its application. *Mutation Research/Reviews in Mutation Research*, 682(1): 71-81.
- Li, W., Zhang, C., Gao, G., Zan, Q., & Yang, Z. (2007). Relationship between Mikania micrantha invasion and soil microbial biomass, respiration and functional diversity. *Plant and Soil*, 296(1-2), 197–207. doi:10.1007/s11104-007-9310-9
- Li, Z.-H., Wang, Q., Ruan, X., Pan, C.-D., & Jiang, D.-A. (2010). *Phenolics and plant allelopathy. Molecules (Basel, Switzerland)*, 15(12), 8933–8952. doi:10.3390/molecules15128933
- Lo Piparo, E., Fratev, F., Lemke, F., Mazzatorta, P., Smiesko, M., Fritz, J. I., & Benfenati, E. (2006). QSAR models for *Daphnia magna* toxicity prediction of benzoxazinone allelochemicals and their transformation products. *Journal of Agricultural and Food Chemistry*, 54(4), 1111–1115. doi:10.1021/jf050918f
- Lorenzo, P, Pazos-Malvido, E., González, L., & Reigosa, M. J. (2008). Allelopathic interference of invasive *Acacia dealbata*: physiological effects. *Allelopathy Journal*, 22(2), 64–76.
- Lorenzo, Paula, González, L., & Reigosa, M. J. (2010). The genus *Acacia* as invader: the characteristic case of *Acacia dealbata* Link in Europe. *Annales Des Sciences Forestières*, 67(1), 101–101. doi:10.1051/forest/2009082
- Lorenzo, Paula, Rodríguez-Echeverría, S., González, L., & Freitas, H. (2010). Effect of invasive *Acacia dealbata* Link on soil microorganisms as determined by PCR-DGGE. *Applied Soil Ecology*, 44(3), 245–251. doi:10.1016/j.apsoil.2010.01.001
- Lowry, O. H., Rosebrough, N. J., Farr, A. L., & Randall, R. J. (1951). Protein measurement with the Folin phenol reagent. *The Journal of Biological Chemistry*, 193(1), 265–275.
- Lu, X. F., Zhang, H., Lyu, S. S., Du, G. D., Wang, X. Q., Wu, C. H., & Lyu, D. G. (2017). Effects of exogenous phenolic acids on photosystem functions and photosynthetic electron transport rate in strawberry leaves. *Photosynthetica*, 56(2), 1–7. doi:10.1007/s11099-017-0702-7
- Mahmood, K. (1998). Effects of salinity, external K⁺/Na⁺ ratio and soil moisture on growth and ion content of *Sesbania rostrata*. *Biologia Plantarum*, 41(2), 297–302.
- Maiti, P., Bhakat, R. K., & Bhattacharjee, A. (2013). Allelopathic potential of a noxious weed on mung bean. *Com Plant Sci*, 3, 31–35.
- Marchante, E., Kjøller, A., Struwe, S., & Freitas, H. (2008). Short- and long-term impacts of *Acacia longifolia* invasion on the belowground processes of a Mediterranean coastal dune ecosystem. *Applied Soil Ecology*, 40(2), 210–217. doi:10.1016/j.apsoil.2008.04.004

- Marimuthu KM, Subramanium K (1960) A haematoxylin squash method for the root tips of *Dolichos lablab* Linn. *Curr. Sci.*, 29: 482-483
- Markham, R., Matthews, R. E. F., & Smith, K. M. (1948). Specific crystalline protein and nucleoprotein from a plant virus having insect vectors. *Nature*, 162(4107), 88-90.
- Markham, R. (1955). Physicochemical studies of the turnip yellow mosaic virus. *Discussions of the Faraday Society*, 11, 221-227.
- Maron, J., & Marler, M. (2007). Native plant diversity resists invasion at both low and high resource levels. *Ecology*, 88(10), 2651–2661. doi:10.1890/06-1993.1
- Marthandan, V., & Jerlin, R. (2017). Effects of Seed Storage Conditions on Biochemical Changes of Freshly Harvested High Moisture Undried Rice Seeds cv. CO 51. *Int. J. Curr. Microbiol. App. Sci.*, 6(12), 2807-2813.
- Materska, M. (2008). Quercetin and its derivatives: chemical structure and bioactivity-a review. *Polish Journal of Food and Nutrition Sciences*, 58(4).
- Matesanz, S., Escudero, A., & Valladares, F. (2008). Additive effects of a potentially invasive grass and water stress on the performance of seedlings of gypsum specialists. *Applied Vegetation Science*, 11(3), 287–296. doi:10.3170/2008-7-18425
- Matthews, E. J., Kruhlak, N. L., Cimino, M. C., Benz, R. D., & Contrera, J. F. (2006). An analysis of genetic toxicity, reproductive and developmental toxicity, and carcinogenicity data: I. Identification of carcinogens using surrogate endpoints. *Regulatory Toxicology and Pharmacology*, 44(2), 83–96. doi:10.1016/j.yrtph.2005.11.003
- McCready, R. M., Guggolz, J., Silviera, V., & Owens, H. S. (1950). Determination of starch and amylose in vegetables. *Analytical Chemistry*, 22(9), 1156–1158. doi:10.1021/ac60045a016
- McPherson, J. K., & Muller, C. H. (1969). Allelopathic Effects of *Adenostoma fasciculatum*, "Chamise", in the California Chaparral. *Ecological Monographs*, 39(2), 177–198. doi:10.2307/1950741
- Mehmood, A., Tanveer, A., Nadeem, M. A., Maqbool, R., & Ali, H. H. (2017). phenolics in two *Alternanthera* species residues affect the germination and early seedling growth of rice (*Oryza sativa*). *Planta Daninha*, 35(0). doi:10.1590/s0100-83582017350100061
- Mersie, W., & Singh, M. (1987). Allelopathic effect of parthenium (*Parthenium hysterophorus* L.) extract and residue on some agronomic crops and weeds. *Journal of Chemical Ecology*, 13(7), 1739–1747. doi:10.1007/BF00980214
- Molisch, H. (1937). Der Einfluss einer Pflanze auf die andere, *Allelopathie*. Fischer Jena.

- Moore, S., & Stein, W. H. (1948). Photometric ninhydrin method for use in the chromatography of amino acids. *Journal of Biological Chemistry*, 176, 367–388
- Moreland, D. E., & Novitzky, W. P. (1987). Effects of phenolic acids, coumarins, and flavonoids on isolated chloroplasts and mitochondria. *Allelochemicals: Role In Agriculture And Forestry*, 330, 247-261..
- Mortelmans, K., & Zeiger, E. (2000). The Ames *Salmonella*/microsome mutagenicity assay. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 455(1-2), 29–60. doi:10.1016/S0027-5107(00)00064-6
- Muller, C. H. (1966). The role of chemical inhibition (allelopathy) in vegetational composition. *Bulletin of the Torrey Botanical Club*, 93(5), 332. doi:10.2307/2483447
- Muller, C. H. (1969). Allelopathy as a factor in ecological process. *Vegetatio Acta Geobotanica*, 18(1-6), 348–357. doi:10.1007/BF00332847
- Navie, S. C., McFadyen, R. E., Panetta, F. D., & Adkins, S. W. (1996). The biology of Australian weeds. 27. *Parthenium hysterophorus* L. *Plant Protection Quarterly*, 11(2), 76–88.
- Nilsson, M.-C., Gallet, C., & Wallstedt, A. (1998). Temporal Variability of Phenolics and Batatasin-III in *Empetrum hermaphroditum* Leaves over an Eight-Year Period: Interpretations of Ecological Function. *Oikos*, 81(1), 6. doi:10.2307/3546462
- Ohno T (2001) *Journal of Environmental Quality* 30: 1631-1635.
- Packialakshmi, N. (n.d.). Alwin. 2014. Analysis of Phytochemical and High Performance Liquid Chromatography in *Desmostachya Bipinnata*. *BMR Biotechnol*, 1, 1–6.
- Pandey, H. N., & Dubey, S. K. (1989). Growth and population dynamics of an exotic weed *Parthenium hysterophorus* Linn. Proceedings: *Plant Sciences*, 99(1), 51–58.
- Pandey, R., Prasad, R. L., Ansari, N. G., & Murthy, R. C. (2015). Utilization of NaOH modified *Desmostachya bipinnata* (Kush grass) leaves and *Bambusa arundinacea* (bamboo) leaves for Cd(II) removal from aqueous solution. *Journal of Environmental Chemical Engineering*, 3(1), 593–602. doi:10.1016/j.jece.2014.06.015
- Panse VG and Sukhatme PV. (1967). *Statistical methods for agricultural workers*. 2nd edition, pp. 150-157. Indian Council of Agriculture Research, New Delhi, India.
- Papadopoulou, K., Melton, R. E., Leggett, M., Daniels, M. J., & Osbourn, A. E. (1999). Compromised disease resistance in saponin-deficient plants. Proceedings of the National

Academy of Sciences of the United States of America, 96(22), 12923–12928.
doi:10.1073/pnas.96.22.12923

Pierik, R., Mommer, L., & Voesenek, L. A. (2013). Molecular mechanisms of plant competition: neighbour detection and response strategies. *Functional Ecology*, 27(4), 841–853.
doi:10.1111/1365-2435.12010

Poitrineau, K., Brown, S. P., & Hochberg, M. E. (2003). Defence against multiple enemies. *Journal of Evolutionary Biology*, 16(6), 1319–1327. doi:10.1046/j.1420-9101.2003.00585.x

Purwar, R. (2019). Antimicrobial textiles. In: *The Impact and Prospects of Green Chemistry for Textile Technology*. pp. 281-306. Elsevier Ltd. Cambridge, MA, USA.

Raizada, P., Raghubanshi, A. S., & Singh, J. S. (2008). *Impact of invasive alien plant species on soil processes: A Review*.

Rajendrudu, G., Prasad, J. S., & Das, V. S. (1986). C(3)-C(4) Intermediate Species in *Alternanthera* (Amaranthaceae): Leaf Anatomy, CO(2) Compensation Point, Net CO(2) Exchange and Activities of Photosynthetic Enzymes. *Plant Physiology*, 80(2), 409–414.
doi:10.1104/pp.80.2.409

Rashmi, S., Singh, S. P., & Deepa, P. (1999). Vegetational profile of macro and microphytes of Jagatdev reservoir. *Environment and Ecology*, 17(3), 674–678.

Reigosa, M. J., Pedrol, N., & González, L. (Eds.). (2006). *Allelopathy*. Dordrecht: Kluwer Academic Publishers. doi:10.1007/1-4020-4280-9

Reinhardt, P., Causà, M., Marian, C. M., & He\|S, B. A. (1996). Adsorption of CO on(110) studied by means of a cluster model surrounded by multipoles obtained from slab calculations. *Physical Review B*, 54(20), 14812–14821. doi:10.1103/PhysRevB.54.14812

Rice E L. 1984. *Allelopathy* pp. 424. 2nd edition. New York, Academic Press. Orlando. FL.

Rizvi, S. J. H., & Rizvi, V. (1992). Exploitation of allelochemicals in improving crop productivity. In S. J. H. Rizvi & V. Rizvi (Eds.), *Allelopathy* (pp. 443–472). Dordrecht: Springer Netherlands. doi:10.1007/978-94-011-2376-1_25

Rodríguez-Echeverría, S. (2009). Organismos del suelo: la dimensión invisible de las invasiones por plantas no nativas. *Revista Ecosistemas*, 18(2).

Romagni, J. G., Rosell, R. C., Nanayakkara, N. P., & Dayan, F. E. (2004). Ecophysiology and potential modes of action for selected lichen secondary metabolites. *Allelopathy*.

- Rout, M. E., & Callaway, R. M. (2009). Plant science. An invasive plant paradox. *Science*, 324(5928), 734–735. doi:10.1126/science.1173651
- Rubaba, O., Chimbari, M., & Mukaratirwa, S. (2017). Scope of research on *Parthenium hysterophorus* in Africa. *South African Journal of Plant and Soil*, 34(5), 323–332. doi:10.1080/02571862.2016.1266046
- Rudrapal, A. B., & Basu, R. N. (1979). Physiology of hydration-dehydration treatment in the maintenance of seed viability in wheat [*Triticum aestivum L.*, India]. *Indian Journal of Experimental Biology*.
- Ruiz, P., Begluitti, G., Tincher, T., Wheeler, J., & Mumtaz, M. (2012). Prediction of acute mammalian toxicity using QSAR methods: a case study of sulfur mustard and its breakdown products. *Molecules (Basel, Switzerland)*, 17(8), 8982–9001. doi:10.3390/molecules17088982
- Sadia Galani. (2011). Germination potential index of Sindh rice cultivars on biochemical basis, using amylase as an indicator. *African Journal of Biotechnology*, 10(80). doi:10.5897/AJB10.1083
- Sage, R. F., Christin, P.-A., & Edwards, E. J. (2011). The C(4) plant lineages of planet Earth. *Journal of Experimental Botany*, 62(9), 3155–3169. doi:10.1093/jxb/err048
- Saini, A., Aggarwal, N. K., Sharma, A., Kaur, M., & Yadav, A. (2014). Utility Potential of *Parthenium hysterophorus* for Its Strategic Management. *Advances in Agriculture*, 2014, 1–16. doi:10.1155/2014/381859
- Sánchez-Del Pino, I., Motley, T. J., & Borsch, T. (2012). Molecular phylogenetics of *Alternanthera gomphrenoideae*, (Amaranthaceae): resolving a complex taxonomic history caused by different interpretations of morphological characters in a lineage with C4 and C3-C4 intermediate species. *Botanical Journal of the Linnean Society*, 169(3), 493–517. doi:10.1111/j.1095-8339.2012.01248.x
- Sánchez-Moreiras, A. M., Martínez-Peña, A., & Reigosa, M. J. (2011). Early senescence induced by 2-3H-benzoxazolinone (BOA) in *Arabidopsis thaliana*. *Journal of Plant Physiology*, 168(9), 863–870. doi:10.1016/j.jplph.2010.11.011
- Sastray, T. C., & Kavathekar, K. Y. (1990). Plants for reclamation of wastelands.
- Saxena, O. P., Singh, G., Pakeeraiah, T., & Pandey, N. (1987). Seed deterioration studies in some vegetable seeds. *Acta Horticulturae*, (215), 39–44. doi:10.17660/ActaHortic.1987.215.5

- Schreiner, O., & Reed, H. S. (1907). The production of deleterious excretions by roots. *Bulletin of the Torrey Botanical Club*, 34(6), 279. doi:10.2307/2479157
- Sekar, K. C. (2012). Invasive alien plants of Indian Himalayan region—Diversity and Implication. *American Journal of Polymer Science*, 03(02), 177–184. doi:10.4236/ajps.2012.32021
- Serafimova, R., Gatnik, M., & Worth, A. (n.d.). Review of QSAR Models and Software Tools for Predicting Genotoxicity and Carcinogenicity.
- Shakila, R., Sri, P. M., Antony, S. A., & Gopakumar, K. (n.d.). Antimicrobial Studies on *Desmostachya bipinnata* Rootstock.
- Shibuya, T. (Agricultural R. C., Tsukuba, Ibaraki (Japan)), Fujii, Y., & Asakawa, Y. (1994). Effects of soil factors on manifestation of allelopathy in *Cytisus scoparius*. *Weed Research* (Japan).
- Singh, A., Saharan, V. A., & Bhandari, A. (2014). Pharmacognostic standardization with various plant parts of *Desmostachya bipinnata*. *Pharmaceutical Biology*, 52(3), 298–307. doi:10.3109/13880209.2013.834367
- Singh, H., Sodhi, N. S., & Singh, N. (2009). Characterisation of starches separated from sorghum cultivars grown in India. *Food Chemistry*, 119(1), 95–100.
- Singh, K. (1994). Site Suitability and Tolerance Limits of Trees, Shrubs and Grasses on Sodic Soils of Ganga-Yamuna Doab. *Indian Forester*.
- Sinha, A., Rana, R. S., & Gupta, S. R. (1991). Growth patterns, net primary production and energy transfers in two grassland communities of sodic soils. *Tropical Ecology*, 32(1), 105–116.
- Smirnoff, N. (1995). Environment and plant metabolism: flexibility and acclimation. *Environment and Plant Metabolism: Flexibility and Acclimation*.
- Soltys, D., Krasuska, U., Bogatek, R., & Gniazdowsk, A. (2013). Allelochemicals as bioherbicides — present and perspectives. In A. Price (Ed.), *Herbicides - Current Research and Case Studies in Use*. InTech. doi:10.5772/56185
- Stickney, J. S., & Hoy, P. R. (1881). Toxic action of black walnut. *Trans. Wis. State Hort. Soc*, 11, 166–167.
- Stoewsand, G. S., Anderson, J. L., Boyd, J. N., Hrazdina, G., Babish, J. G., Walsh, K. M., & Losco, P. (1984). Quercetin: A mutagen, not a carcinogen, in Fischer rats. *Journal of Toxicology and Environmental Health*, 14(2-3), 105–114. doi:10.1080/15287398409530565

- Sullivan, K. M., Manuppello, J. R., & Willett, C. E. (2014). Building on a solid foundation: SAR and QSAR as a fundamental strategy to reduce animal testing. *SAR and QSAR in Environmental Research*, 25(5), 357–365. doi:10.1080/1062936X.2014.907203
- Tamado, T., Schutz, W., & Milberg, P. (2002). Germination ecology of the weed *Parthenium hysterophorus* in eastern Ethiopia. *Annals of Applied Biology*, 140(3), 263–270. doi:10.1111/j.1744-7348.2002.tb00180.x
- Tamaro, M., Monti-Bragadin, C., & Banfi, E. (1975). Mutagenic activity of anthraquinone derivatives used as dyes in a textile factory. *Bollettino dell'Istituto Sieroterapico Milanese*, 54(2), 105–107.
- Taye, T., Gossman, M., Einhorn, G., Büttner, C., Metz, R., & Abate, D. (2002). The potential of pathogens as biological control of parthenium weed (*Parthenium hysterophorus L.*) in Ethiopia. *Mededelingen (Rijksuniversiteit Te Gent. Fakulteit van de Landbouwkundige En Toegepaste Biologische Wetenschappen)*, 67(3), 409–420.
- Tharayil, N. (2009). To survive or to slay: Resource-foraging role of metabolites implicated in allelopathy. *Plant Signaling & Behavior*, 4(7), 580–583. doi:10.4161/psb.4.7.8915
- Trethewey, R. N., & Smith, A. M. (2000). Starch metabolism in leaves. In *Photosynthesis* (pp. 205–231). Springer, Dordrecht.
- Van der Putten, W. H., Van Dijk, C., & Peters, B. A. M. (1993). Plant-specific soil-borne diseases contribute to succession in foredune vegetation. *Nature*, 362(6415), 53–56. doi:10.1038/362053a0
- Van der WAL, R., Truscott, A.-M., Pearce, I. S. K., Cole, L., Harris, M. P., & Wanless, S. (2008). Multiple anthropogenic changes cause biodiversity loss through plant invasion. *Global Change Biology*, 14(6), 1428–1436. doi:10.1111/j.1365-2486.2008.01576.x
- Van Wilgen, B. W., Reyers, B., Le Maitre, D. C., Richardson, D. M., & Schonegevel, L. (2008). A biome-scale assessment of the impact of invasive alien plants on ecosystem services in South Africa. *Journal of Environmental Management*, 89(4), 336–349. doi:10.1016/j.jenvman.2007.06.015
- Vartak, V. D. (1968). Weed that threatens crop and grasslands in Maharashtra. *Indian Farming*, 18(1).
- Venkataiah, B., Ramesh, C., Ravindranath, N., & Das, B. (2003). Charminarone, a seco-pseudoguaianolide from *Parthenium hysterophorus*. *Phytochemistry*, 63(4), 383–386.

- Vilà, M., Basnou, C., Gollasch, S., Josefsson, M., Pergl, J., & Scalera, R. (2009). One hundred of the most invasive alien species in europe. *A Handbook of Alien Species In Europe* (pp. 265–268). Dordrecht: Springer Netherlands. doi:10.1007/978-1-4020-8280-1_12
- Walpole, M., Almond, R. E. A., Besançon, C., Butchart, S. H. M., Campbell-Lendrum, D., Carr, G. M., Zimsky, M. (2009). Ecology. Tracking progress toward the 2010 biodiversity target and beyond. *Science*, 325(5947), 1503–1504. doi:10.1126/science.1175466
- Wardle, D. A., Bardgett, R. D., Klironomos, J. N., Setälä, H., van der Putten, W. H., & Wall, D. H. (2004). Ecological linkages between aboveground and belowground biota. *Science*, 304(5677), 1629–1633. doi:10.1126/science.1094875
- Weber, H., Heim, U., Golombek, S., Borisjuk, L., & Wobus, U. (1998). Assimilate uptake and the regulation of seed development. *Seed Science Research*, 8(3), 331-346.
- Weidenhamer, J. D. (1996). Distinguishing resource competition and chemical interference: overcoming the methodological impasse. *Agronomy Journal*, 88(6), 866. doi:10.2134/agronj1996.00021962003600060005x
- Whittaker, R. H., & Feeny, P. P. (1971). Allelochemistry: chemical interactions between species. *Science*, 171(3973), 757–770. doi:10.1126/science.171.3973.757
- Wigglesworth, V. B. (1952). The role of iron in histological staining. *Journal of Cell Science*, 3(21), 105–118.
- Williams, J. D., & Groves, R. H. (1980). The influence of temperature and photoperiod on growth and development of *Parthenium hysterophorus L.* *Weed Research*, 20(1), 47–52. doi:10.1111/j.1365-3180.1980.tb00040.x
- Wink, M. (2003). Evolution of secondary metabolites from an ecological and molecular phylogenetic perspective. *Phytochemistry*, 64(1), 3–19. doi:10.1016/S0031-9422(03)00300-5
- Wittmann, W. (1962). Aceto-iron-haematoxylin for staining chromosomes in squashes of plant material. *Stain Technology*, 37, 27–30. doi:10.3109/10520296209114565
- Wolfreys, A. M., & Hepburn, P. A. (2002). Safety evaluation of phytosterol esters. Part 7. *Assessment of mutagenic activity of phytosterols, phytosterol esters and the cholesterol derivative, 4-cholest-3-one*. *Food and Chemical Toxicology*, 40(4), 461–470. doi:10.1016/s0278-6915(01)00099-0
- Wu, H., Haig, T., Pratley, J., Lemerle, D., & An, M. (2002). Biochemical basis for wheat seedling allelopathy on the suppression of annual ryegrass (*Lolium rigidum*). *Journal of Agricultural and Food Chemistry*, 50(16), 4567–4571. doi:10.1021/jf025508v

- Yang, R. Z., & Tang, C. S. (1988). Plants used for pest control in China: A literature review. *Economic Botany*, 42(3), 376–406. doi:10.1007/BF02860162
- Zackrisson, O., & Nilsson, M.-C. (1992). Allelopathic effects by *Empetrum hermaphroditum* on seed germination of two boreal tree species. *Canadian Journal of Forest Research*, 22(9), 1310–1319. doi:10.1139/x92-174
- Zhang, C. B., Wang, J., Qian, B. Y., & Li, W. H. (2009). Effects of the invader *Solidago canadensis* on soil properties. *Applied Soil Ecology*, 43(2-3), 163–169. doi:10.1016/j.apsoil.2009.07.001
- Zhou, Y. H., & Yu, J. Q. (2006). Allelochemicals and photosynthesis. In M. J. Reigosa, N. Pedrol, & L. González (Eds.), *Allelopathy* (pp. 127–139). Dordrecht: Kluwer Academic Publishers. doi:10.1007/1-4020-4280-9_6
- Zhu, H., Martin, T. M., Ye, L., Sedykh, A., Young, D. M., & Tropsha, A. (2009). Quantitative structure-activity relationship modeling of rat acute toxicity by oral exposure. *Chemical Research in Toxicology*, 22(12), 1913–1921. doi:10.1021/tx900189p
- Zobel, A. M., Lynch, J. M., & Jeffrey, M. (1997). Extrusion of UV-absorbing phenolics in Acer spp. In response to UV and freezing temperature. I. UV A-absorbing compounds on the surface of *Acer saccharum* and *Acer platanoides* autumn leaves. *Allel. J*, 4, 269–276.