

LIST OF FIGURES

| | | |
|------------|--|----|
| Fig. 1.1. | Schematic representation of ZnO wurtzite crystal structure. | 7 |
| Fig. 1.2. | Transitions ascribed to various defect levels in ZnO. | 9 |
| Fig. 1.3. | Schematic presentation of magnetic polaron model. | 15 |
| Fig. 1.4. | Schematic image of magnetic polarons in a semiconductor lattice. | 16 |
| Fig. 2.1. | Chemical precipitation method. | 26 |
| Fig. 2.2. | A presentation of Bragg's diffraction of X-ray beam from atomic planes. | 29 |
| Fig. 2.3. | Representation of XRD experimental set-up. | 30 |
| Fig. 2.4. | A schematic view of FTIR spectrophotometer. | 33 |
| Fig. 2.5. | Working principle of FTIR spectrophotometer. | 33 |
| Fig. 2.6. | Schematic set-up for absorption measurements. | 35 |
| Fig. 2.7. | Schematic set-up of Lambda-35 double beam UV/Vis spectrophotometer. | 35 |
| Fig. 2.8. | The schematic of fluorescence experimental set-up. | 36 |
| Fig. 2.9. | Photo degradation over length. | 38 |
| Fig. 3.1. | EDX spectra of (a) Zn _{0.94} Ni _{0.06} O and (b) Zn _{0.91} Ni _{0.09} O samples. | 45 |
| Fig. 3.2. | XRD spectra of Zn _{1-x} Ni _x O. | 46 |
| Fig. 3.3. | TEM image of (a) Zn _{0.94} Ni _{0.06} O and (b) Zn _{0.91} Ni _{0.09} O samples. | 50 |
| Fig. 3.4. | HRTEM image of (a) Zn _{0.94} Ni _{0.06} O and (b) Zn _{0.91} Ni _{0.09} O samples. | 50 |
| Fig. 3.5. | SAED pattern of (a) Zn _{0.94} Ni _{0.06} O and (b) Zn _{0.91} Ni _{0.09} O samples. | 51 |
| Fig. 3.6. | FTIR spectra of Zn _{1-x} Ni _x O series for (a) 400-700 cm ⁻¹ and (b) 700-4000 cm ⁻¹ . | 52 |
| Fig. 3.7. | Absorption spectra of Zn _{1-x} Ni _x O series. | 53 |
| Fig. 3.8. | FL spectra of Zn _{1-x} Ni _x O series. | 54 |
| Fig. 3.9. | Nyquist plot of (a) pure ZnO, (b) Zn _{0.97} Ni _{0.03} O and (c) Zn _{0.91} Ni _{0.09} O samples. | 56 |
| Fig. 3.10. | Variation of (a) dielectric constant and (b) loss tangent with frequency of Ni-doped ZnO. | 57 |
| Fig. 3.11. | P-E loop of Zn _{0.91} Ni _{0.09} O and Zn _{0.85} Ni _{0.15} O sample. | 58 |
| Fig. 3.12. | M-H loop of Zn _{0.91} Ni _{0.09} O sample by SQUID. | 59 |

| | | |
|------------|--|----|
| Fig. 3.13. | Variation of ME voltage coefficient with DC magnetic field of $\text{Zn}_{0.91}\text{Ni}_{0.09}\text{O}$ and $\text{Zn}_{0.85}\text{Ni}_{0.15}\text{O}$ samples. | 60 |
| Fig. 4.1. | XRD patterns of $\text{Zn}_{1-x}\text{Co}_x\text{O}$ series. | 66 |
| Fig. 4.2. | (a) TEM image and (b) HRTEM photograph of $\text{Zn}_{0.94}\text{Co}_{0.06}\text{O}$ sample. | 67 |
| Fig. 4.3. | (a) SAED pattern and (b) EDX spectra of $\text{Zn}_{0.94}\text{Co}_{0.06}\text{O}$ sample. | 67 |
| Fig. 4.4. | FTIR spectra of $\text{Zn}_{1-x}\text{Co}_x\text{O}$ series for (a) 400-800 cm^{-1} and (b) 700-4000 cm^{-1} . | 69 |
| Fig. 4.5. | Absorption spectra of $\text{Zn}_{1-x}\text{Co}_x\text{O}$ series. | 70 |
| Fig. 4.6. | FL spectra of $\text{Zn}_{1-x}\text{Co}_x\text{O}$ series. | 71 |
| Fig. 4.7. | Variation of (a) dielectric constant and (b) loss with frequency of Co-doped ZnO nanoparticles. | 72 |
| Fig. 4.8. | M-H curve of $\text{Zn}_{0.94}\text{Co}_{0.06}\text{O}$ sample by SQUID. | 73 |
| Fig. 4.9. | Variation of ME voltage coefficient with DC magnetic field of $\text{Zn}_{0.82}\text{Co}_{0.18}\text{O}$ sample. | 74 |
| Fig. 5.1. | XRD patterns of $\text{Zn}_{1-x}\text{Fe}_x\text{O}$ series. | 82 |
| Fig. 5.2. | Rietveld refinement of XRD patterns of $\text{Zn}_{1-x}\text{Fe}_x\text{O}$ for $x = 0, 0.03, 0.06$ and 0.09 . | 82 |
| Fig. 5.3. | TEM image of (a) $\text{Zn}_{0.97}\text{Fe}_{0.03}\text{O}$ and (b) $\text{Zn}_{0.85}\text{Fe}_{0.15}\text{O}$ samples. | 84 |
| Fig. 5.4. | HRTEM image of (a) $\text{Zn}_{0.97}\text{Fe}_{0.03}\text{O}$ sample and (b) $\text{Zn}_{0.85}\text{Fe}_{0.15}\text{O}$ sample. | 84 |
| Fig. 5.5. | SAED pattern of (a) $\text{Zn}_{0.97}\text{Fe}_{0.03}\text{O}$ and (b) $\text{Zn}_{0.85}\text{Fe}_{0.15}\text{O}$ samples. | 84 |
| Fig. 5.6. | EDX spectra of $\text{Zn}_{0.97}\text{Fe}_{0.03}\text{O}$ sample. | 85 |
| Fig. 5.7. | FTIR spectra of $\text{Zn}_{1-x}\text{Fe}_x\text{O}$ series (400 cm^{-1} - 5000 cm^{-1}). | 86 |
| Fig. 5.8. | Absorption spectra of $\text{Zn}_{1-x}\text{Fe}_x\text{O}$ series. | 87 |
| Fig. 5.9. | $(\alpha h\nu)^2$ vs. $h\nu$ plot of $\text{Zn}_{1-x}\text{Fe}_x\text{O}$ series. | 88 |
| Fig. 5.10. | FL spectra of $\text{Zn}_{1-x}\text{Fe}_x\text{O}$ series. | 88 |
| Fig. 5.11. | Variation of (a) dielectric constant and (b) loss with frequency of $\text{Zn}_{1-x}\text{Fe}_x\text{O}$ series. | 90 |
| Fig. 5.12. | P-E loop of undoped ZnO, $\text{Zn}_{0.94}\text{Fe}_{0.06}\text{O}$, $\text{Zn}_{0.91}\text{Fe}_{0.09}\text{O}$ and $\text{Zn}_{0.88}\text{Fe}_{0.12}\text{O}$ samples. | 91 |
| Fig. 5.13. | M-H loop of $\text{Zn}_{1-x}\text{Fe}_x\text{O}$ sample. | 93 |

| | | |
|------------|---|---------|
| Fig. 5.14. | ABK plot of $Zn_{0.97}Fe_{0.03}O$ sample. | 93 |
| Fig. 6.1. | Photocatalytic set-up for MB degradation by Mg-doped ZnO. | 99 |
| Fig. 6.2. | XRD patterns of $Zn_{1-x}Mg_xO$ samples. | 102 |
| Fig. 6.3. | TEM image of (a) $Zn_{0.94}Mg_{0.06}O$, (b) $Zn_{0.88}Mg_{0.12}O$ and (c) $Zn_{0.85}Mg_{0.15}O$ samples. | 104 |
| Fig. 6.4. | HRTEM image of $Zn_{0.85}Mg_{0.15}O$ sample. | 104 |
| Fig. 6.5. | SAED pattern of (a) $Zn_{0.88}Mg_{0.12}O$ and (b) $Zn_{0.85}Mg_{0.15}O$ samples. | 104 |
| Fig. 6.6. | EDX pattern of $Zn_{0.85}Mg_{0.15}O$ sample. | 105 |
| Fig. 6.7. | FTIR spectra of $Zn_{1-x}Mg_xO$ series ($400\text{ cm}^{-1} - 4000\text{ cm}^{-1}$). | 105 |
| Fig. 6.8. | (a) Absorption spectra of $Zn_{1-x}Mg_xO$ series and (b) Current density vs. electric field graph. | 107 |
| Fig. 6.9. | FL spectra of $Zn_{1-x}Mg_xO$ series. | 108 |
| Fig. 6.10. | Photocatalytic degradation of (a) pure ZnO, (b) $Zn_{0.97}Mg_{0.03}O$, (c) $Zn_{0.94}Mg_{0.06}O$, (d) $Zn_{0.91}Mg_{0.09}O$, (e) $Zn_{0.88}Mg_{0.12}O$ and (f) $Zn_{0.85}Mg_{0.15}O$. | 110-111 |
| Fig. 6.11. | Kinetic study of MB degradation by $Zn_{1-x}Mg_xO$. | 111 |
| Fig. 6.12. | Variation of (a) ϵ and (b) $\tan\delta$ with frequency and composition of $Zn_{1-x}Mg_xO$. | 112 |
| Fig. 6.13. | P-E loop of $Zn_{0.91}Mg_{0.09}O$ & $Zn_{0.85}Mg_{0.15}O$ samples. | 113 |
| Fig. 6.14. | M-H loop of $Zn_{0.85}Mg_{0.15}O$ sample. | 114 |
| Fig. 6.15. | ME loop of (a) $Zn_{0.91}Mg_{0.09}O$ and (b) $Zn_{0.88}Mg_{0.12}O$ samples. | 115 |
| Fig. 7.1. | Photocatalytic set-up for MB degradation by Nd-doped ZnO. | 120 |
| Fig. 7.2. | XRD patterns of $Zn_{1-x}Nd_xO$ series. | 121 |
| Fig. 7.3. | (a) TEM image, (b) HRTEM image, (c) SAED pattern, (d) EDX pattern of $Zn_{0.91}Nd_{0.09}O$ sample. | 123 |
| Fig. 7.4. | Absorption spectra of $Zn_{1-x}Nd_xO$ series. | 124 |
| Fig. 7.5. | FL spectra of $Zn_{1-x}Nd_xO$ series. | 125 |
| Fig. 7.6. | Photocatalytic degradation of (a) pure ZnO (b) $Zn_{0.97}Nd_{0.03}O$ (c) $Zn_{0.94}Nd_{0.06}O$ (d) $Zn_{0.91}Nd_{0.09}O$ (e) $Zn_{0.88}Nd_{0.12}O$ and (f) $Zn_{0.85}Nd_{0.15}O$. | 128 |
| Fig. 7.7. | Kinetic study of methylene blue degradation. | 129 |