
List of Figures

| Chapter-1. Introduction: | Page-no. |
|--|-----------------|
| Fig.1. Heart Coronary arteries | 3 |
| Fig.2. Atherosclerotic plaque in coronary arteries | 4 |
| Fig.3. Different heart valves | 7 |
| Fig.4. Heart cardiomyopathy | 8 |
| Fig.5. Flow chart of different types of cardiovascular diseases | 10 |
| Chapter-2. Review and Literatures: | |
| Fig.1. Epidemiology of disease mortality | 17 |
| Fig.2. The stages of development of an atherosclerotic plaque | 21 |
| Fig.3. Development of atherosclerotic plaque with superimposed thrombus | 22 |
| Fig.4. Role of insulin for disease pathogenesis | 26 |
| Fig.5. Intraabdominal fat contributes to insulin resistance and cardiovascular dysfunction through cytokine | 30 |
| Chapter-5. Result: | |
| Fig.1. Follow-up of the body weight changes of rats | 48 |
| Fig.2. Catalase activity and superoxide dismutase activity (SOD1) of rat liver & heart tissues by gel-zymography | 49 |
| Fig.3. Protein band pattern in PAGE from serum sample of cardiac patients | 51 |
| Fig.4. Nutritional status (BMI) wise distribution and dispersion pattern | 54 |
| Fig.5. Dermcidin (DCN-2) induced synthesis of TNF- α | 55 |
| Fig.6. Synthesis of IL-6 in neutrophils by DCN-2 | 56 |

| | |
|--|----|
| Fig.7. Identification of dermcidin in the cell free plasma of both normal and type-1 diabetes mellitus patients. | 57 |
| Fig.8. Bar diagram shows comparison of blood glucose, uric acid, cholesterol and triglyceride of male Wister rats | 58 |
| 6. Discussions: | |
| Fig.9. Superoxide dismutase activity in human serum | 63 |
| Fig.10. Bar diagram shows comparison of MDA (a) and NPSH (b) level in blood serum of male & female cardiac patients | 66 |
| Fig.11. Nutritional status wise distribution patterns (%) of individuals from different groups of blood glucose level | 69 |
| Fig.12. Comparison by Student's t test of underweight and overweight blood glucose levels | 70 |
| Fig.13. Estriol effect to inhibit TNF- α level in patients with coronary artery disease (CAD) neutrophils. | 72 |
| Fig.14. Role of estriol on IL-6 level in CAD patients. | 73 |
| Fig.15. Plasma level of C-reactive protein, malondialdehyde and non-protein soluble thiol in T1DM and normal subjects and DCN-2 induced synthesis of TNF- α in liver cells: | 78 |
| Fig.16. Bar diagram shows the blood serum hs CRP & TSH level comparison among patients with cardiac risk factors & cardiac | 81 |
| Fig.17. Bar diagram shows the blood serum T3 & T4 level comparison among cardiac risk factors & cardiac marker | 83 |

| | |
|--|----|
| Fig.18. Plasma level of NO and insulin in normal and T1DM volunteers and their correlation with plasma dermcidin level | 84 |
| Fig 19. DCN-2 induced inhibition of estriol mediated NO synthesis inneutrophils. Different concentrations of DCN-2 | 86 |
| Fig.20. CRP regulating mechanistic pathway on coronary artery disease via inflammatory and coagulation process. | 87 |
| Fig.21. Role of oxidative stress on vascular damage followed by inflammatory responses via excess ROS production. | 88 |
